

8632

Access DB# 81051

## SEARCH REQUEST FORM

Scientific and Technical Information Center

Requester's Full Name: Kedechi, Ekwine Examiner #: 77000 Date: 11/25/02  
 Art Unit: 1213 Phone Number 30 6-570 / Serial Number: 10/010883  
 Mail Box and Bldg/Room Location: 323 9A11 Results Format Preferred (circle): PAPER DISK E-MAIL

If more than one search is submitted, please prioritize searches in order of need.

\*\*\*\*\*  
 Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc, if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.


Title of Invention: Stabilization of aqueous emulsion polymer  
 Inventors (please provide full names): Natale et al.

Earliest Priority Filing Date: 7/99

\*For Sequence Searches Only\* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.

Please search for the reaction products in claim 26 in a polymer emulsion.

The possible reaction products are:



## STAFF USE ONLY

Searcher: <u>Ed</u>	Type of Search	Vendors and cost where applicable
Searcher Phone #: _____	NA Sequence (#) _____	STN <u>\$ 547.29</u>
Searcher Location: _____	AA Sequence (#) _____	Dialog _____
Date Searcher Picked Up: _____	Structure (#) <u>✓ (5) (abstracts)</u>	Questel/Orbit _____
Date Completed: <u>12-5-02</u>	Bibliographic <u>✓ (and)</u>	Dr. Link _____
Searcher Prep & Review Time: <u>10</u>	Litigation _____	Lexis/Nexis _____
Clerical Prep Time: _____	Fulltext _____	Sequence Systems _____
Online Time: <u>115</u>	Patent Family _____	WWW/Internet _____
	Other _____	Other (specify) _____

BEST AVAILABLE COPY

=> file reg

FILE 'REGISTRY' ENTERED AT 10:53:43 ON 05 DEC 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 3 DEC 2002 HIGHEST RN 475040-20-1  
DICTIONARY FILE UPDATES: 3 DEC 2002 HIGHEST RN 475040-20-1

TSCA INFORMATION NOW CURRENT THROUGH MAY 20, 2002

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. See HELP  
PROPERTIES for more information. See STNote 27, Searching Properties  
in the CAS Registry File, for complete details:  
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=> d his

(FILE 'HOME' ENTERED AT 09:22:33 ON 05 DEC 2002)

FILE 'HCAPLUS' ENTERED AT 09:25:51 ON 05 DEC 2002

L1 557 S NATALE ?/AU  
L2 635 S KLIMA ?/AU  
L3 5846 S HOPKINS ?/AU  
L4 1239 S WIGGINS ?/AU  
L5 1 S L1 AND L2 AND L3 AND L4  
SEL L5 1 RN

FILE 'REGISTRY' ENTERED AT 09:26:13 ON 05 DEC 2002

L6 5 S E1-E5  
L7 1 S 106-89-8  
L8 21605 S 106-89-8/CRN  
SEL L6 1,2,4 RN  
L9 3 S E6-E8  
EDIT E6-E8 /BI /CRN  
L10 135 S E6-E8  
L11 0 S L8 AND L10

FILE 'HCAPLUS' ENTERED AT 09:30:05 ON 05 DEC 2002

L12 13239 S L7  
L13 1085 S L9  
L14 16 S L12 AND L13  
L15 101612 S (EMULS? OR MICROEMULS? OR DISPERS? OR MICRODISPERS? OR  
L16 4 S L14 AND L15

L17 170564 S (AQ# OR AQUEOUS? OR WATER? OR H2O) (2A) (EMULS? OR MICROE  
L18 3 S L14 AND L17  
L19 46751 S (STABIL? OR STABL?) (2A) (EMULS? OR MICROEMULS? OR DISPER  
L20 1 S L14 AND L19

FILE 'LREGISTRY' ENTERED AT 09:37:17 ON 05 DEC 2002

L21 STR  
L22 STR  
L23 STR  
L24 STR

FILE 'REGISTRY' ENTERED AT 09:48:12 ON 05 DEC 2002

L25 SCR 2043  
L26 2 S L21 AND L25  
L27 SCR 963 AND 1700 AND 1707  
L28 7 S L21 AND L25 AND L27  
L29 SCR 963 AND 1700  
L30 20 S (L21 OR (L22 AND (L23 OR L24))) AND L25 AND L29  
L31 3215 S (L21 OR (L22 AND (L23 OR L24))) AND L25 AND L29 FUL  
SAV L31 EGW883/A

FILE 'LREGISTRY' ENTERED AT 09:54:55 ON 05 DEC 2002

L32 STR

FILE 'REGISTRY' ENTERED AT 09:58:16 ON 05 DEC 2002

L33 4 S L32 SSS SAM SUB=L31  
L34 STR L32  
L35 0 S L34 SSS SAM SUB=L31  
L36 0 S L34 SSS FUL SUB=L31  
L37 319 S L31 AND 1/NC  
E OXIRANE/CN  
L38 1 S E3  
L39 21344 S 75-21-8/CRN  
E METHYLOXIRANE/CN  
L40 1 S E3  
L41 17754 S 75-56-9/CRN  
L42 8 S L31 AND (L39 OR L41) AND 2/NC  
L43 190 S L31 AND L39 AND L41 AND 3/NC  
L44 198 S L42 OR L43

FILE 'LREGISTRY' ENTERED AT 10:09:55 ON 05 DEC 2002

L45 226 S (C H CL/ELF OR C H BR/ELF OR C H I/ELF) AND 2<X

FILE 'LCA' ENTERED AT 10:10:13 ON 05 DEC 2002

L46 0 S ALKANETRIYL?

FILE 'HCAPLUS' ENTERED AT 10:10:25 ON 05 DEC 2002

L47 63 S ALKANETRIYL?

FILE 'LREGISTRY' ENTERED AT 10:12:09 ON 05 DEC 2002

FILE 'REGISTRY' ENTERED AT 10:12:41 ON 05 DEC 2002

L48 21318 S (C H CL/ELF OR C H BR/ELF OR C H I/ELF) AND 2<X  
 L49 0 S L48 AND L31  
 L50 19563 S L48 NOT PMS/CI  
 L51 14457 S L50 NOT M/ELS  
 L52 11225 S L51 AND 1/NC

FILE 'HCAPLUS' ENTERED AT 10:15:29 ON 05 DEC 2002

L53 101482 S L52  
 L54 12752 S L37  
 L55 1815 S L44  
 L56 129 S L53 AND (L54 OR L55)  
 L57 12889 S L52 (L) RCT/RL  
 L58 385 S L37 (L) RCT/RL  
 L59 48 S L44 (L) RCT/RL  
 L60 1 S L57 AND (L58 OR L59)  
 L61 18 S L56 AND (L15 OR L17 OR L19)  
 L62 42842 S (BASE# OR BASIC?) (2A) (CAT# OR CATALY?)  
 L63 0 S L56 AND L62  
 L64 11835 S (NAOH OR KOH OR OH OR HYDROXIDE#) (2A) (CAT# OR CATALY?)  
 L65 7861 S WILLIAMSON# OR ETHER# (2A) (SYN# OR SYNTH?)  
 L66 0 S L56 AND (L64 OR L65)  
 L67 QUE REACT? OR RXN#  
 L68 30 S L56 AND L67  
 L69 6 S L16 OR L18 OR L20 OR L60  
 L70 18 S L61 NOT L69  
 L71 24 S L68 NOT (L69 OR L70)  
 L72 11 S L14 NOT (L69 OR L70 OR L71)

FILE 'REGISTRY' ENTERED AT 10:53:43 ON 05 DEC 2002

=> d l31 que stat

L21 STR

Ak^G1—OH O~^Ak  
 1 2 3 @6 @7

REP G1=(1-10) 6-1 7-3

NODE ATTRIBUTES:

CONNECT IS E1 RC AT 1  
 CONNECT IS E2 RC AT 7  
 DEFAULT MLEVEL IS ATOM  
 GGCAT IS SAT AT 1  
 GGCAT IS SAT AT 7  
 DEFAULT ECLEVEL IS LIMITED  
 ECOUNT IS M3 C AT 1  
 ECOUNT IS M2-X3 C AT 7

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
 NUMBER OF NODES IS 5

STEREO ATTRIBUTES: NONE



L22 STR

Ak—OH

1 2

## NODE ATTRIBUTES:


CONNECT IS E1 RC AT 1  
 DEFAULT MLEVEL IS ATOM  
 GGCAT IS SAT AT 1  
 DEFAULT ECLEVEL IS LIMITED  
 ECOUNT IS M3 C AT 1

## GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
 NUMBER OF NODES IS 2

## STEREO ATTRIBUTES: NONE

L23 STR



## NODE ATTRIBUTES:


DEFAULT MLEVEL IS ATOM  
 DEFAULT ECLEVEL IS LIMITED

## GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
 NUMBER OF NODES IS 3

## STEREO ATTRIBUTES: NONE

L24 STR



## NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM  
 DEFAULT ECLEVEL IS LIMITED

## GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
 NUMBER OF NODES IS 4

## STEREO ATTRIBUTES: NONE

L25 SCR 2043

L29 SCR 963 AND 1700  
L31 3215 SEA FILE=REGISTRY SSS FUL (L21 OR (L22 AND (L23 OR  
L24))) AND L25 AND L29

100.0% PROCESSED 228865 ITERATIONS  
SEARCH TIME: 00.00.07

3215 ANSWERS

=> file hcaplus  
FILE 'HCAPLUS' ENTERED AT 10:56:38 ON 05 DEC 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 5 Dec 2002 VOL 137 ISS 23  
FILE LAST UPDATED: 3 Dec 2002 (20021203/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

CAS roles have been modified effective December 16, 2001. Please check your SDI profiles to see if they need to be revised. For information on CAS roles, enter HELP ROLES at an arrow prompt or use the CAS Roles thesaurus (/RL field) in this file.

=> d l69 1-6 cbib abs hitstr hitind


L69 ANSWER 1 OF 6 HCAPLUS COPYRIGHT 2002 ACS  
2002:237363 Document No. 136:263994 **Stabilization of aqueous emulsion polymers.** Natale, Marcie; Klima, Rudolph; Hopkins, Thomas; Wiggins, Michael (Cognis Corporation, USA). U.S. US 6362259 B1 20020326, 9 pp. (English). CODEN: USXXAM. APPLICATION: US 1999-363555 19990729.  
AB An **emulsion polymer** compn. contains an **emulsion stabilizing** effective quantity of product consisting essentially of the base catalyzed reaction of (A) an epihalohydrin and (B) at least one compd. R(OE)n(OP)mOH, wherein R is a satd. or unsatd. org. group having from 3 to 22 carbon atoms. n is a no. of from 1 to 50, m is a no. from 0 to 10, EO represents an ethyleneoxy group, and OP represents a propyleneoxy group, wherein

the mole ratio of components (A) to component (B) is from about 0.60:1 to about 2:1. A reaction product of epichlorohydrin and ethoxylated decyl alc. was used to stabilize a Bu acrylate-vinyl acetate copolymer.

IT 106-89-8DP, Epichlorohydrin, reaction products with alkoxyated alcs. 9004-77-7DP, Polyethylene glycol monobutyl ether, reaction products with epichlorohydrin 26183-52-8DP, Ethoxylated decyl alcohol, reaction products with epichlorohydrin 27252-75-1DP, Polyethylene glycol monooctyl ether, reaction products with epichlorohydrin (stabilization of aq. emulsion polymers)


RN 106-89-8 HCAPLUS

CN Oxirane, (chloromethyl) - (9CI) (CA INDEX NAME)




RN 9004-77-7 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-butyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)




RN 26183-52-8 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



RN 27252-75-1 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-octyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IC ICM C08K005-06  
ICS C08K005-1515  
NCL 524114000  
CC 37-6 (Plastics Manufacture and Processing)  
ST **aq emulsion polymer stabilization**  
IT **Emulsifying agents**  
**Emulsions**  
**(stabilization of aq. emulsion polymers)**  
IT 25067-01-0P, Butyl Acrylate-Vinyl Acetate copolymer  
**(stabilization of aq. emulsion polymers)**  
IT 106-89-8DP, Epichlorohydrin, reaction products with alkoxyated alcs. 9004-77-7DP, Polyethylene glycol monobutyl ether, reaction products with epichlorohydrin 26183-52-8DP, Ethoxylated decyl alcohol, reaction products with epichlorohydrin 27252-75-1DP, Polyethylene glycol monooctyl ether, reaction products with epichlorohydrin  
**(stabilization of aq. emulsion polymers)**


L69 ANSWER 2 OF 6 HCAPLUS COPYRIGHT 2002 ACS  
2001:676845 Document No. 135:228259 Branched polymer defoamers for pigment dispersants. Breindel, Kenneth; Brown, David W. (Cognis Corporation, USA). PCT Int. Appl. WO 2001066629 A1 20010913, 21 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2001-US7552 20010309. PRIORITY: US 2000-PV188091 20000309; US 2001-801509 20010308.

AB The compn., useful in water-based coatings, comprises (A) a pigment dispersant; and (B) a branched polymer defoamer obtained by reaction of (a) a linking agent RY<sub>3</sub>, wherein each Y is a halogen atom or one Y is halogen atom and two Y groups together represent an epoxy oxygen atom, which is attached to two adjacent carbon atoms in the R group to form an epoxy group, and R is C<sub>3</sub>-10 alkanetriyl, and (b) .gtoreq.1 compd. R<sub>1</sub>X(EO)<sub>n</sub>(PO)<sub>m</sub>(BO)<sub>p</sub>Z (R<sub>1</sub> = (un)substituted satd. or unsatd., C<sub>4</sub>-36 org. group; X = O, S, NR<sub>2</sub>; R<sub>2</sub> = H, C<sub>1</sub>-4 alkyl; Z = H, NHR<sub>2</sub>, SH; n = 0-100; m = 0-50; p = 0-50; n + m + p .gtoreq.1; EO = residue of ethylene oxide; PO = residue of propylene oxide; BO = residue of butylene oxide). Thus, 100 mL 1% aq. soln. of Tamol 731 (diisobutylene-maleic anhydride **copolymer pigment dispersant**) was mixed with 1 drop of reaction product of epichlorohydrin and decyl alc. 4EO in 1:1 epoxy : OH ratio to give a clear soln., 1 g of which was added to 99 mL water and hand shaken, showing no stable foam.

IT 106-89-8DP, Epichlorohydrin, reaction products with polyethylene glycol monodecyl ether 26183-52-8DP, Polyethylene glycol monodecyl ether, reaction products with epichlorohydrin (defoamers; branched polymer defoamers for pigment dispersants)


RN 106-89-8 HCAPLUS

CN Oxirane, (chloromethyl)- (9CI) (CA INDEX NAME)



RN 26183-52-8 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IC ICM C08K005-06

CC 42-5 (Coatings, Inks, and Related Products)

ST branched **polymer** defoamer pigment **dispersant** coating; epichlorohydrin polyoxyethylene decyl ether reaction defoamer; diisobutylene maleic anhydride **copolymer** pigment **dispersant**

IT 106-89-8DP, Epichlorohydrin, reaction products with polyethylene glycol monodecyl ether 9002-92-0DP, Polyethylene glycol monodecyl ether, reaction products with epichlorohydrin 9016-45-9P, Polyethylene glycol monononylphenol ether 26183-52-8DP, Polyethylene glycol monodecyl ether, reaction products with epichlorohydrin (defoamers; branched polymer defoamers for pigment dispersants)

IT 31051-58-8, Maleic anhydride-methacrylic acid-styrene **copolymer** 37199-81-8, TAMOL 731 (pigment **dispersants**; branched **polymer** defoamers for pigment dispersants)

L69 ANSWER 3 OF 6 HCAPLUS COPYRIGHT 2002 ACS

2001:25785 Document No. 134:87944 Demulsifiers for separating oil and water mixtures. Lindert, Andreas; Wiggins, Michael S. (Henkel Corporation, USA). U.S. US 6172123 B1 20010109, 5 pp. (English). CODEN: USXXAM. APPLICATION: US 1999-364417 19990730.


AB A method for preventing or breaking oil and H<sub>2</sub>O mixts. comprises adding to emulsion a branched reaction product of (A) linking compds. R1X3 (R1 = C3-10 alkanetriyl; X = halo, or 1 X group = halo and 2 X groups with 2 adjacent C atoms in R1 and an O atom form an

epoxy group), preferably epichlorohydrin, (B) (alkoxylated) alcs. R2(OA)nOH (R2 = C3-36 alkyl; OA = CH2CH2O, CH2CHMeO, CH2CHEtO; n = 1-100), and/or (alkoxylated) aliph. amines R2(OA)nNH2 (R2, OA, n as above), and, optionally other polyols and polyamines. The branched products are low-foaming surfactants which are highly effective when used alone as demulsifiers esp. in preventing or breaking the interlayer present between oil and H2O layers obtained in the recovery of petroleum crude oil from underground sources. For example, adding 0.01% of an epichlorohydrin reaction product with ethoxylated (4 EO) decyl alc. to aq. diesel fuel **emulsion** caused sepn. into 2 distinct layers within 1 min.

IT **106-89-8D**, Epichlorohydrin, reaction products with ethoxylated decyl alc. **26183-52-8D**, Polyethylene glycol decyl ether, reaction products with epichlorohydrin (low-foam surfactants as demulsifiers for sepg. oil and water mixts.)


RN 106-89-8 HCAPLUS

CN Oxirane, (chloromethyl)- (9CI) (CA INDEX NAME)



RN 26183-52-8 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IC ICM B01D017-05

NCL 516179000

CC 46-4 (Surface Active Agents and Detergents)

ST oil **water emulsion** sepn low foam surfactant demulsifier; diesel oil emulsion breaking ethoxylated decyl alc epichlorohydrin product; epichlorohydrin ethoxylated decyl alc product demulsifier aq diesel oil


IT **106-89-8D**, Epichlorohydrin, reaction products with ethoxylated decyl alc. **26183-52-8D**, Polyethylene glycol decyl ether, reaction products with epichlorohydrin (low-foam surfactants as demulsifiers for sepg. oil and water mixts.)

L69 ANSWER 4 OF 6 HCAPLUS COPYRIGHT 2002 ACS


1999:763948 Document No. 132:23832 Antistatic lubricant composition and method of making same. Wiggins, Michael; Incorvia, Michael J.;

Fischer, Stephen A. (Henkel Corporation, USA). PCT Int. Appl. WO 9961169 A1 19991202, 26 pp. DESIGNATED STATES: W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2. APPLICATION: WO 1999-US10458 19990525. PRIORITY: US 1998-85134 19980526.


- AB A process for providing a substrate with antistatic and lubricating properties by contacting the substrate with a water-sol. or **water-dispersible polymer** compn. having hydrophobic and hydrophilic properties. The compn. comprises polyethers obtained from ethoxylated or ethoxylated-propoxylated fatty alcs. having a terminal hydrocarbon chain length of at least one carbon.
- IT **26183-52-8**, Polyethylene glycol decyl ether  
(Trycol 5950; ethoxylated and propoxylated fatty alc.-based antistatic lubricant compn. for textiles and plastics and method of making same)
- RN 26183-52-8 HCAPLUS
- CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)




- IT **106-89-8**, reactions **9004-77-7**, Polyethylene glycol monobutyl ether  
(ethoxylated and propoxylated fatty alc.-based antistatic lubricant compn. for textiles and plastics and method of making same)
- RN 106-89-8 HCAPLUS
- CN Oxirane, (chloromethyl)- (9CI) (CA INDEX NAME)



- RN 9004-77-7 HCAPLUS
- CN Poly(oxy-1,2-ethanediyl), .alpha.-butyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)




- IC ICM B05D003-12  
ICS B32B027-00; B32B027-06; B32B027-32; B32B027-36  
CC 40-9 (Textiles and Fibers)  
Section cross-reference(s): 38  
IT 26183-52-8, Polyethylene glycol decyl ether  
(Trycol 5950; ethoxylated and propoxylated fatty alc.-based antistatic lubricant compn. for textiles and plastics and method of making same)  
IT 106-89-8, reactions 9004-77-7, Polyethylene glycol monobutyl ether  
(ethoxylated and propoxylated fatty alc.-based antistatic lubricant compn. for textiles and plastics and method of making same)
- L69 ANSWER 5 OF 6 HCAPLUS COPYRIGHT 2002 ACS  
1976:4596 Document No. 84:4596 Applications of surfactants to synthetic organic chemistry. Menger, F. M.; Rhee, J. U.; Rhee, H. K. (Dep. Chem., Emory Univ., Atlanta, Ga., USA). J. Org. Chem., 40(25), 3803-5 (English) 1975. CODEN: JOCEAH.
- AB Surfactant additives can give higher yields and shorter reactions times in reactions of water-insol. org. liqs. with aq. reagents. Piperonal was oxidized to piperonylic acid and PhCCl<sub>3</sub> was hydrolyzed to benzoic acid ion media contg. quaternary ammonium bromide and Brij 35.
- IT 98-07-7  
(hydrolysis of, surfactants as solubilization agents in)
- RN 98-07-7 HCAPLUS  
CN Benzene, (trichloromethyl)- (9CI) (CA INDEX NAME)



- IT 9002-92-0  
(solubilization agent for oxidn. of piperonal and hydrolysis of benzylidyne trichloride)
- RN 9002-92-0 HCAPLUS  
CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)





CC 25-17 (Noncondensed Aromatic Compounds)

Section cross-reference(s): 46

IT 98-07-7

(hydrolysis of, surfactants as solubilization agents in)

IT 57-09-0 1643-19-2 9002-92-0

(solubilization agent for oxidn. of piperonal and hydrolysis of benzylidene trichloride)

L69 ANSWER 6 OF 6 HCAPLUS COPYRIGHT 2002 ACS

1974:427077 Document No. 81:27077 Acrylonitrile copolymer. Sato, Hideo; Ueda, Takeshi (Asahi Chemical Industry Co., Ltd.). Japan. JP 48033993 B4 19731018 Showa, 9 pp. (Japanese). CODEN: JAXXAD. APPLICATION: JP 1970-74540 19700827.

AB Antistatic fibers are prepd. by spinning copolymers of acrylonitrile with  $\text{CH}_2:\text{C}(\text{R}_1)\text{CO}_2\text{CH}_2\text{CH}(\text{OH})\text{CH}_2(\text{OCH}_2\text{CH}_2)_n\text{OR}_2$  (I,  $\text{R}_1 = \text{H}, \text{Me}; \text{R}_2 = \text{n-alkyl}$ ) or  $\text{CH}_2:\text{C}(\text{R}_3)\text{CO}_2\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{O}_2\text{CZCO}(\text{OCH}_2\text{CH}_2)_n\text{R}_4$  (II,  $\text{R}_3 = \text{H}, \text{Me}; \text{R}_4 = \text{Cl}, \text{CN}, \text{alkoxy}, \text{or alkanoylamino}; \text{Z} = (\text{CH}_2)_4, \text{CH}_2\text{CH}_2, \text{or o-C}_6\text{H}_4$ ). Thus, 1 mole polyethylene glycol mono-n-octyl ether [27252-75-1] (d.p. 15) was condensed with 1.05 moles epichlorohydrin [106-89-8] in the presence of  $\text{BF}_3 \cdot \text{OEt}_2$  and the product was treated either directly as the chlorohydrin ether or after cyclization to the glycidyl ether, with acrylic acid [79-10-7] to give the vinyl monomer I ( $\text{R}_1 = \text{Me}; \text{R}_2 = \text{n-C}_8\text{H}_{17}, n = 15$ ) (III) [51877-41-9]. Suspension polym. of III with acrylonitrile with a redox catalyst gave an 82.5:17.5 acrylonitrile-polyethylene glycol 3-acryloyloxy-2-hydroxypropyl n-octyl ether copolymer [51937-37-2], reduced viscosity (0.3g/100 ml DMF, 35.deg.) 1.26. A soln. of 27 parts of this copolymer in 140 parts 70%  $\text{HNO}_3$  at 0.deg. was spun into 33%  $\text{HNO}_3$  at 0.deg. to give fibers with surface resistance 4 .tim. 1010 .OMEGA. (7 .tim. 1010 .OMEGA. after washing) and dry strength 3.0 g/denier, compared with 5 .tim. 1013 .OMEGA. and 3.1 g/denier for 90.5:9.0:0.5 acrylonitrile-methyl acrylate-sodium allylsulfonate copolymer (IV) [25053-78-5] fibers. Similar fibers were prepd. from 5 other acrylonitrile-I or -II copolymers. A 65:35 copolymer of acrylonitrile with II [ $\text{R}_3 = \text{Me}, \text{R}_4 = \text{CN}, \text{Z} = (\text{CH}_2)_4, n = 10$ ] was spun in a 15:85 blend with IV to give fibers with surface resistance 5 .tim. 1010 .OMEGA. before and 6 .tim. 1010 .OMEGA. after washing. Five similar blends of acrylonitrile-I or -II copolymers with IV were similarly spun. A 15:85 copolymer of acrylonitrile with II ( $\text{R}_3 = \text{R}_4 = \text{Me}, \text{Z} = \text{CH}_2\text{CH}_2, n = 25$ ) was prepd. and used in 1.5% aq. soln. as an antistatic finish for acrylic fiber garments.


IT 27252-75-1P

(prepn. of)

RN 27252-75-1 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-octyl-.omega.-hydroxy- (9CI) (CA

INDEX NAME)




IT 27252-75-1

(reaction of, with epichlorohydrin)

RN 27252-75-1 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-octyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)




IT 106-89-8, reactions

(with polyethylene glycol monoethyl ether)

RN 106-89-8 HCAPLUS

CN Oxirane, (chloromethyl)- (9CI) (CA INDEX NAME)



IC C08F; D01F

CC 39-2 (Textiles)

IT 27252-75-1P 52641-09-5P 52656-36-7P  
(prepn. of)

IT 27252-75-1

(reaction of, with epichlorohydrin)

IT 106-89-8, reactions

(with polyethylene glycol monoethyl ether)

=&gt; d l72 1-11 ti

L72 ANSWER 1 OF 11 HCAPLUS COPYRIGHT 2002 ACS

TI Defoamers for aqueous systems such as latex paints

L72 ANSWER 2 OF 11 HCAPLUS COPYRIGHT 2002 ACS

TI Defoamers for aqueous systems such as latex paints

L72 ANSWER 3 OF 11 HCAPLUS COPYRIGHT 2002 ACS

TI Polymeric lubricant compositions, and methods for their use

L72 ANSWER 4 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
TI Gloss retention additives for cleaning compositions

L72 ANSWER 5 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
TI Alkyl polyglycoside compositions having reduced viscosity and inhibited crystallization

L72 ANSWER 6 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
TI Water bath and method for electrolytic deposition of copper coatings

L72 ANSWER 7 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
TI Ion-conducting polymer compositions showing temperature-independent conductivity

L72 ANSWER 8 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
TI Polyoxyalkene substituted and bridged triazine, benzotriazole and benzophenone derivatives as UV absorbers

L72 ANSWER 9 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
TI Preparation of glycidyl ethers from alcohols and epichlorohydrin

L72 ANSWER 10 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
TI Modified polyamines


L72 ANSWER 11 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
TI Amphoteric surfactants

=> d l72 2,3,11 cbib abs hitstr hitind


L72 ANSWER 2 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
2002:10328 Document No. 136:71294 Defoamers for aqueous systems such as latex paints. Wiggins, Michael S.; Broadbent, Ronald W. (Cognis Corp., USA). PCT Int. Appl. WO 2002000319 A1 20020103, 23 pp.  
DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English).  
CODEN: PIXXD2. APPLICATION: WO 2001-US20360 20010626. PRIORITY: US 2000-606092 20000627.

AB The products of the reaction of epichlorohydrin and compds. having the formula (I)  $R_3(EO)_n(PO)_mOH$  wherein  $R_3$  is an alkyl, alkenyl or arenyl group having from 4 to 22 carbon atoms; a substituted alkyl or alkenyl group having from 4 to 22 carbon atoms wherein;  $n$  is a no. from 0 to 50 and  $m$  is a no. from 0 to 50; wherein the mole ratio of epichlorohydrin to I is from .apprx.0.60/1 to .apprx.2/1 are used in defoaming compns. for defoaming aq. systems such as latex paints.


IT 106-89-8, Epichlorohydrin, reactions 9004-77-7D,  
 Butanol, ethoxylated 26183-52-8, Trycol 5950  
 26183-52-8D, Decyl alcohol, ethoxylated 27252-75-1D  
 , Octyl, ethoxylated  
 (defoamers for aq. systems such as latex paints)  
 RN 106-89-8 HCAPLUS  
 CN Oxirane, (chloromethyl)- (9CI) (CA INDEX NAME)




RN 9004-77-7 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-butyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)




RN 26183-52-8 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



RN 26183-52-8 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



RN 27252-75-1 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-octyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IC ICM B01D019-04  
 ICS B01F017-42; B01F017-56; C11D001-72; C11D003-22  
 CC 42-5 (Coatings, Inks, and Related Products)  
 IT 106-89-8, Epichlorohydrin, reactions 108-88-3, Toluene, reactions 109-63-7, Boron trifluoride etherate 124-41-4, Sodium methoxide 1310-73-2, Sodium hydroxide, reactions 9004-77-7D, Butanol, ethoxylated 26183-52-8, Trycol 5950 26183-52-8D, Decyl alcohol, ethoxylated 27252-75-1D, Octyl, ethoxylated (defoamers for aq. systems such as latex paints)


L72 ANSWER 3 OF 11 HCAPLUS COPYRIGHT 2002 ACS  
 2001:360117 Document No. 134:369235 Polymeric lubricant compositions, and methods for their use. Wiggins, Michael S.; Incorvia, Michael J.; Klima, Rudolph F.; Boudreaux, Chase J. (Cognis Corporation, USA). PCT Int. Appl. WO 2001034736 A1 20010517, 34 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2.  
 APPLICATION: WO 2000-US28750 20001018. PRIORITY: US 1999-PV161445 19991026; US 2000-672218 20000928.

AB The method of the invention comprises reducing friction and wear between solid surfaces which are, or which are to be, in relative motion, comprising applying to at least one of the solid surfaces a polymeric material as a synthetic lubricating oil. The lubricating oils can be used to lubricate solid surfaces where elevated operating temps. are present.

IT 106-89-8, Epichlorohydrin, reactions 9004-77-7 26183-52-8 27252-75-1 (polymeric lubricant compns., and methods for their use)


RN 106-89-8 HCAPLUS

CN Oxirane, (chloromethyl)- (9CI) (CA INDEX NAME)




RN 9004-77-7 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-butyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)




RN 26183-52-8 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



RN 27252-75-1 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-octyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IC ICM C10M107-20

ICS C10M107-40

CC 51-8 (Fossil Fuels, Derivatives, and Related Products)

IT 67-56-1, Methanol, reactions 79-11-8, Chloroacetic acid, reactions 106-89-8, Epichlorohydrin, reactions 112-70-9D, Tridecyl alcohol, ethoxylated 124-41-4, Sodium methoxide 624-48-6, Dimethyl maleate 1310-73-2, Sodium hydroxide, reactions 7681-57-4, Sodium metabisulfite 9004-77-7 9016-45-9 24938-91-8 25322-68-3, Polyethylene glycol 26183-52-8 27252-75-1 29797-40-8 65605-36-9, Jeffamine ED-6000 (polymeric lubricant compns., and methods for their use)


L72 ANSWER 11 OF 11 HCAPLUS COPYRIGHT 2002 ACS

1985:455716 Document No. 103:55716 Amphoteric surfactants. (Toho Chemical Industry Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP 60019030 A2 19850131 Showa, 5 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 1983-126188 19830713.


AB Surfactants RO(ZO)<sub>n</sub>CH<sub>2</sub>CHOHCH<sub>2</sub>N+HR<sub>1</sub>CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub><sup>-</sup> (I; R = C<sub>8</sub>-24 alkyl, alkenyl; R<sub>1</sub> = C<sub>1</sub> to req. 5 alkyl; Z = ethylene, ethylene and propylene; n = 1-50) are prepd. by reaction of 1 mol epihalohydrin with 1 mol polyoxyalkylene alkyl (or alkenyl) ether, followed by treatment with 1 mol C<sub>1</sub> to req. 5 alkylamine and alkali to replace the terminal halogen with NHR<sub>1</sub>, then addn. reaction with a lower alkyl

acrylate, hydrolysis, and neutralization with acid. Thus, 1 mol epichlorohydrin and 1 mol polyoxyethylene lauryl ether (d.p. 5) were mixed 8 h at 70.degree. with 0.5 g Et<sub>2</sub>O.BF<sub>3</sub> to obtain 1 mol 1-chloro-2-hydroxy-3-(lauryloxypolyethyleneoxy)propane [97332-31-5], which was mixed with 1 mol MeNH<sub>2</sub> [74-89-5] and heated 4 h at 120.degree. to obtain N-[lauryloxy(polyethyleneoxy)hydroxypropyl]-N-methylamine (II) [97332-32-6] (amine no. 113.8). Then, 0.5 mol II and 0.5 mol Me acrylate [96-33-3] were mixed and heated at 120.degree. for 10 h, treated with an NaOH soln. at 70.degree. for 10 h, then neutralized to obtain 272 g N-[lauryloxy(polyethyleneoxy)hydroxypropyl]-N-methylalanine [96743-35-0], which was sol. in water, EtOH, 1% aq. NaOH, and 1% aq. HCl, and whose 1% and 0.1% aq. solns. both showed surface tension 32 dyne/cm.

IT 26183-52-8DP, reaction products with epihalohydrins, amines and acrylate esters, hydrolyzed  
(oligomeric, amphoteric surfactants, prepn. and properties of)  
RN 26183-52-8 HCAPLUS  
CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IT 106-89-8, reactions  
(reaction of, with polyoxyalkylene alkyl ethers)  
RN 106-89-8 HCAPLUS  
CN Oxirane, (chloromethyl)- (9CI) (CA INDEX NAME)



IC ICM B01F017-42  
ICS B01F017-52  
CC 46-3 (Surface Active Agents and Detergents)  
IT 9002-92-0DP, reaction products with epihalohydrins, amines and acrylate esters, hydrolyzed 9004-95-9DP, reaction products with epihalohydrins, amines and acrylate esters, hydrolyzed 9004-98-2DP, reaction products with epihalohydrins, amines and acrylate esters, hydrolyzed 9005-00-9DP, reaction products with epihalohydrins, amines and acrylate esters, hydrolyzed 24938-91-8DP, reaction products with epihalohydrins, amines and acrylate esters, hydrolyzed 26183-52-8DP, reaction products with epihalohydrins, amines and acrylate esters, hydrolyzed 37311-00-5DP, reaction products with epihalohydrins, amines and

acrylate esters, hydrolyzed 37311-04-9DP, reaction products with ethylamine, Et 2-propenoate and bromomethyloxirane, hydrolyzed (oligomeric, amphoteric surfactants, prepn. and properties of)  
IT 106-89-8, reactions 3132-64-7  
(reaction of, with polyoxyalkylene alkyl ethers)

=> d l70 1-18 cbib abs hitstr hitind

L70 ANSWER 1 OF 18 HCAPLUS COPYRIGHT 2002 ACS

2001:427377 Document No. 135:15441 Adjuvants for herbicidal compositions, providing enhanced effectiveness. Brinker, Ronald J.; Dyszlewski, Andrew D.; Gillespie, Jane L.; Jones, Claude R.; Kramer, Richard M.; Pallas, Norman R.; Radke, Rodney O.; Ward, Anthony J. I.; Xu, Xiaodong C. (Monsanto Company, USA). U.S. US 6245713 B1 20010612, 40 pp., Cont.-in-part of U.S. 6,184,182. (English). CODEN: USXXAM. APPLICATION: US 1999-298136 19990423. PRIORITY: US 1996-PV29317 19961025; US 1997-PV34887 19970131; US 1997-PV39789 19970304; US 1997-957750 19971024.


AB A compn. comprises, dissolved or **dispersed** in **water**, an anionic exogenous chem. substance such as the herbicide glyphosate, together with: (i) alkyl ether surfactant(s)  $[R1O(CH_2CH_2O)_n[(CHR)_{20}]mR_2]$  R1 = aliph. satd. or unsatd. C16-22 hydrocarbyl; n = 5-100; m = 0,1-5; R = H, Me or CHR<sub>2</sub>O; R<sub>2</sub> = H, C1-4 alkyl or C2-4 acyl]; and (ii) amine surfactant(s) each having a mol. structure that comprises (a) a hydrophobic moiety having one or a plurality of independently satd. or unsatd., branched or unbranched, aliph., alicyclic or arom. C3-20 hydrocarbyl or hydrocarbylene groups joined together by 0 to about 7 ether linkages and having in total about 8 to about 24 carbon atoms, and (b) a hydrophilic moiety comprising an amino group that is cationic or that can be protonated to become cationic, having attached directly thereto 1 to 3 oxyethylene groups or polyoxyethylene chains, these oxyethylene groups and polyoxyethylene chains comprising on av. 1 to about 50 oxyethylene units per surfactant mol., the hydrophobic moiety being attached either to the amino group or via an ether linkage to an oxyethylene unit. The wt. ratio of the alkyl ether surfactant(s) to the amine surfactant(s) is about 1:10 to about 10:1; and the alkyl ether and amine surfactants are present in total in an adjuvant amt. of about 0.05 to about 0.5 parts by wt. per part by wt. of the herbicide, expressed as acid equiv. Also provided are solid and liq. conc. compns. that can be dild., dissolved or **dispersed** in **water** to form such a plant treatment compn.

IT 71-55-6  
(adjuvants for herbicidal compns., providing enhanced effectiveness)


RN 71-55-6 HCAPLUS

CN Ethane, 1,1,1-trichloro- (8CI, 9CI) (CA INDEX NAME)






IT 9002-92-0, laureth-4 9005-00-9, brij 78  
 (adjuvants for herbicidal compns., providing enhanced effectiveness)  
 RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



RN 9005-00-9 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-octadecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



IC ICM A01N025-30  
 ICS A01N057-02  
 NCL 504206000  
 CC 5-3 (Agrochemical Bioregulators)  
 IT 50-31-7, 2,3,6-TBA 71-55-6 75-99-0, Dalapon 85-34-7,  
 Fenac 93-65-2, Mecoprop 94-74-6, MCPA 94-75-7, 2,4-D,  
 biological studies 94-81-5, MCPB 94-82-6, 2,4-DB 112-05-0,  
 Nonanoic acid 120-36-5, Dichlorprop 124-58-3, Methylarsonic acid  
 132-66-1, Naptalam 133-90-4, Chloramben 145-73-3, Endothall  
 314-40-9, Bromacil 1071-83-6, Glyphosate 1689-83-4, Ioxynil  
 1689-84-5, Bromoxynil 1702-17-6, Clopyralid 1918-00-9, Dicamba  
 1918-02-1, Picloram 3337-71-1, Asulam 3813-05-6, Benazolin  
 5329-14-6, Sulfamic acid 25057-89-0, Bentazon 35597-43-4,  
 Bilanafos 38641-94-0, Roundup 40465-66-5, ammonium glyphosate  
 40843-25-2, Diclofop 50594-66-6, Acifluorfen 51276-47-2,  
 Glufosinate 55335-06-3, Triclopyr 58667-63-3, Flamprop  
 59682-52-9, Fosamine 69335-91-7, Fluazifop 69806-34-4, Haloxyfop  
 72178-02-0, Fomesafen 76578-12-6, Quizalofop 77501-60-1,  
 Fluoroglycofen 81334-34-1, Imazapyr 81335-37-7, Imazaquin  
 81335-77-5, Imazethapyr 84087-01-4, Quinclorac 87547-04-4,  
 Flumiclorac 95617-09-7, Fenoxaprop 100728-84-5, Imazamethabenz

104098-48-8, Imazameth 114311-32-9, Imazamox  
(adjuvants for herbicidal compns., providing enhanced effectiveness)

IT 9002-92-0, laureth-4 9005-00-9, brij 78  
(adjuvants for herbicidal compns., providing enhanced effectiveness)

L70 ANSWER 2 OF 18 HCAPLUS COPYRIGHT 2002 ACS


2000:674636 Document No. 133:224212 Fluorine-based oil and water repellent composition having better stability in storage. Lee, Soo-bok; Kim, Dong-kwon; Lee, Kwang-won; Kim, Kwang-je; Park, Inn-joon (Korea Research Institute of Chemical Technology, S. Korea). Repub. Korea KR 9700319 B1 19970108, No pp. given (Korean). CODEN: KRXXFC. APPLICATION: KR 1994-37129 19941227.

AB The mixt. of perfluoroalkylethyl acrylate, stearyl methacrylate, vinylidene chloride, N-methylolacrylamide, polyoxyethylene lauryl ether, stearyltrimethylammonium chloride, acetate, distd. water, azobisisobuthylamidine bihydrochloric acid salt and 1,1,2-trichloroethylene, are fed into the pressure reactor which is attached with a temp. adjusting device and an agitator, and which is made of a glass. The mixt. is stirred up and nitrogen substitution is made for 30 min. The temp. of the reactive soln. is risen to 60.degree. and polymer processing is made for 15 h, cooled off and a **water dispersion** type fluorine group water repelling agent in which 20% solid substance is included is produced.

IT 79-01-6, 1,1,2-Trichloroethylene, uses 9002-92-0, Polyoxyethylene lauryl ether  
(compn. contg.; prepn. of fluorine-based oil and water repellent compn. having better stability in storage)


RN 79-01-6 HCAPLUS

CN Ethene, trichloro- (9CI) (CA INDEX NAME)



RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



IC ICM D06M011-07

CC 40-9 (Textiles and Fibers)

IT 79-01-6, 1,1,2-Trichloroethylene, uses 112-03-8,

Stearyltrimethylammonium chloride 9002-92-0,  
Polyoxyethylene lauryl ether

(compn. contg.; prepn. of fluorine-based oil and water repellent  
compn. having better stability in storage)

L70 ANSWER 3 OF 18 HCAPLUS COPYRIGHT 2002 ACS

1999:723543 Document No. 131:326866 Development and application of  
microemulsion for the in situ extraction of polychlorinated  
biphenyls from soil. Dierkes, Frank (Inst. Chem. Dynamik Geosphere  
7 Angewandte Physikalische Chem., Forschungszentrum Juelich G.m.b.H.,  
Juelich, D-52425, Germany). Berichte des Forschungszentrums Juelich,  
Juel-3679, 1-131 pp. (German) 1999. CODEN: FJBEE5. ISSN:  
0366-0885.


AB For the extn. of polychlorinated biphenyls a microemulsion  
consisting of biol. decomposable components is developed. The  
**microemulsion** remains **stable** at .apprx.10.degree.  
and is usable in the in situ soil remediation. The microemulsion  
system consists of equal amts. of water (with addn. of CaCl<sub>2</sub>) and  
rape oil Me ester. Na-bis(2-ethylhexyl)sulfosuccinate (AOT) and  
2,5,8-trimethylnonylhexaglycoether (Tergitol TMN 6) are used as  
amphiphilic compds. At low temps. the undesired formation of liq.  
crystals in the microemulsion can be suppressed by increasing the  
electrolyte concn. in the water phase, but this leads to an  
unfavorable shift of the phase boundaries of the system. The  
substitution of a part of the nonionic surfactant by an ethoxylated  
castor oil suppresses the formation of liq. crystals. The extn.  
performance of the microemulsion system is investigated with doped  
soils in batch and glass column expts. The contaminant is removed  
nearly completely from the arable soil. Extn. rate is lower on  
sandy soil. The microemulsion system can be splitted specifically  
into an oil phase and a water phase by increasing the temp. From  
the tech. point of view the reuse of the water phase is uncrit., but  
legislative problems can arise.

IT 7012-37-5, PCB 28 35065-27-1, PCB 153  
35065-28-2, PCB 138 35065-29-3, PCB 180  
35693-99-3, PCB 52 37680-73-2, PCB 101

(development and application of microemulsion for in situ extn.  
of polychlorinated biphenyls from soil)


RN 7012-37-5 HCAPLUS

CN 1,1'-Biphenyl, 2,4,4'-trichloro- (9CI) (CA INDEX NAME)




RN 35065-27-1 HCAPLUS

CN 1,1'-Biphenyl, 2,2',4,4',5,5'-hexachloro- (9CI) (CA INDEX NAME)




RN 35065-28-2 HCAPLUS

CN 1,1'-Biphenyl, 2,2',3,4,4',5'-hexachloro- (9CI) (CA INDEX NAME)




RN 35065-29-3 HCAPLUS

CN 1,1'-Biphenyl, 2,2',3,4,4',5,5'-heptachloro- (9CI) (CA INDEX NAME)




RN 35693-99-3 HCAPLUS

CN 1,1'-Biphenyl, 2,2',5,5'-tetrachloro- (9CI) (CA INDEX NAME)



RN 37680-73-2 HCAPLUS

CN 1,1'-Biphenyl, 2,2',4,5,5'-pentachloro- (9CI) (CA INDEX NAME)




IT 60828-78-6, Tergitol TMN6

(development and application of microemulsion for in situ extn. of polychlorinated biphenyls from soil)

RN 60828-78-6 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-[3,5-dimethyl-1-(2-methylpropyl)hexyl]-.omega.-hydroxy- (9CI) (CA INDEX NAME)

CC 60-4 (Waste Treatment and Disposal)  
Section cross-reference(s): 19IT 118-74-1, HCB 7012-37-5, PCB 28 35065-27-1, PCB  
153 35065-28-2, PCB 138 35065-29-3, PCB 180  
35693-99-3, PCB 52 37680-73-2, PCB 101

(development and application of microemulsion for in situ extn. of polychlorinated biphenyls from soil)

IT 577-11-7 60828-78-6, Tergitol TMN6 192391-56-3, RME

(development and application of microemulsion for in situ extn. of polychlorinated biphenyls from soil)


L70 ANSWER 4 OF 18 HCAPLUS COPYRIGHT 2002 ACS

1999:455757 Document No. 131:134138 Single-phase microemulsification of a complex light-nonaqueous-phase-liquid: laboratory evaluation of several mixtures of surfactant/alcohol solutions. Rhue, R. Dean; Suresh, P.; Rao, C.; Annable, Michael D. (Soil and Water Science Dep., Gainesville, FL, 32611, USA). Journal of Environmental Quality, 28(4), 1135-1144 (English) 1999. CODEN: JEVQAA. ISSN: 0047-2425. Publisher: American Society of Agronomy.


AB A recent advance in conventional pump-and-treat technol. for aquifer remediation involves the use of surfactant-alc. mixts. that will form a clear, transparent, thermodynamically stable oil-in-**water microemulsion** on contact with a residual non-aq.-phase-liq. (NAPL). An initial screening of 86 com.-grade

surfactants for aq. soly. resulted in selection of 58 that were further tested in batch expts. to evaluate the capacity to solubilize a complex NAPL waste collected from a Superfund site (Operable Unit OU-1) at Hill AFB, UT. The selected group of 58 surfactants represented 6 classes of anionic, 9 classes of nonionic, and one class of amphoteric surfactants. Batch studies on NAPL solubilization identified a no. of surfactants suitable for use in the field demonstration phase of the project; a further criterion in surfactant selection was that the flushing soln. had a viscosity <2 cp. The best surfactants among this group had HLB (hydrophilic-lipophilic balance) values 12-13, and solubilized 10-20 g/L of the OU-1 NAPL when the surfactant concn. was 3%. Column tests using NAPL-coated glass beads showed that the more efficient surfactants could remove >90% of the NAPL after flushing with <10 pore vols. Brij 97, an ethoxylated alc. ether surfactant, showed a high capacity for solubilizing the OU-1 NAPL. In a column test using contaminated Hill AFB aquifer material, flushing with a mixt. of 3% Brij 97 and 2.5% n-pentanol removed essentially all of the mass of 9 target analytes in the NAPL after flushing with <10 pore vols. without mobilizing the NAPL or destabilizing aquifer colloids.

IT 9002-92-0, Macol LA 790  
 (Brij 30, Brij 35, Macol LA 790; single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)  
 RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




IT 9004-95-9  
 (Brij 56, Brij 58; single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)  
 RN 9004-95-9 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




IT 24938-91-8, Hetoxol TD 9  
 (Hetoxol TD 9; single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)

RN 24938-91-8 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-tridecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




IT 26183-52-8, Trycol 5953  
 (Trycol 5953; single-phase microemulsification of complex  
 light-nonaq.-phase-liq. and evaluation of mixts. of  
 surfactant/alc. solns.)

RN 26183-52-8 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA  
 INDEX NAME)




IT 9002-92-0 9005-00-9, Brij 78  
 (single-phase microemulsification of complex light-nonaq.-phase-  
 liq. and evaluation of mixts. of surfactant/alc. solns.)

RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




RN 9005-00-9 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-octadecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



IT 120-82-1, 1,2,4-Trichlorobenzene  
 (single-phase microemulsification of complex light-nonaq.-phase-  
 liq. and evaluation of mixts. of surfactant/alc. solns.)

RN 120-82-1 HCAPLUS  
 CN Benzene, 1,2,4-trichloro- (8CI, 9CI) (CA INDEX NAME)



- CC 61-2 (Water)  
Section cross-reference(s): 46, 60
- IT **Water** purification  
(**microemulsification**; single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)
- IT 9002-92-0, Macol LA 790  
(Brij 30, Brij 35, Macol LA 790; single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)
- IT 9004-95-9  
(Brij 56, Brij 58; single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)
- IT 24938-91-8, Hetoxol TD 9  
(Hetoxol TD 9; single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)
- IT 26183-52-8, Trycol 5953  
(Trycol 5953; single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)
- IT 71-41-0, n-Pentanol, uses 151-21-3, Sodium dodecyl sulfate, uses 922-80-5, Aerosol AY 65 2373-38-8, Aerosol MA 80I  
9002-92-0 9002-92-0 9002-92-0  
9002-92-0 9002-92-0 9004-82-4, Witcolate S-1285C  
9004-99-3, Myrj-52 9005-00-9, Brij 78 9005-65-6, Tween 80 9005-66-7, Tween 40 9005-67-8, Tween 60 27731-62-0, Standapol ES-40 60371-17-7, Antarox LF-330 72175-39-4, Glucamate SSE-20 123898-54-4, Witcolate SE-5 138673-02-6, Alkasurf LA-EP 15 182915-72-6, Alcodet SK 205132-39-4, Antarox LF-224 234758-46-4, Emkapon Jel BS 234763-43-0, Syn Fac TDA 92 234764-72-8, Abitec WA 664 234767-37-4, Antarox LA-EP 25LF 234767-38-5, Antarox LA-EP 45 234767-41-0, Rexonic P 5 234767-46-5, Drewpol 10-1CCK 234768-21-9, DeSulf GOS-P 70  
(single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)
- IT 91-20-3, Naphthalene, processes 95-50-1, 1,2-Dichlorobenzene 95-63-6, 1,2,4-Trimethylbenzene 112-40-3, n-Dodecane 120-82-1, 1,2,4-Trichlorobenzene 124-18-5, n-Decane



629-50-5, n-Tridecane 1120-21-4, n-Undecane 1330-20-7, Xylene, processes  
(single-phase microemulsification of complex light-nonaq.-phase-liq. and evaluation of mixts. of surfactant/alc. solns.)

L70 ANSWER 5 OF 18 HCAPLUS COPYRIGHT 2002 ACS


1999:163464 Document No. 130:227036 Analytical monitoring of photocatalytic treatments. Degradation of 2,3,6-trichlorobenzoic acid in **aqueous TiO2 dispersions**. Prevot, Alessandra Bianco; Pramauro, Edmondo (Dipartimento di Chimica Analitica, Universita di Torino, Turin, 10125, Italy). Talanta, 48(4), 847-857 (English) 1999. CODEN: TLNTA2. ISSN: 0039-9140. Publisher: Elsevier Science B.V..

AB Photocatalytic degrdn. of 2,3,6-trichlorobenzoic acid (2,3,6-TBA) in **aq. TiO2 dispersions** irradiated with simulated solar light was studied. Fast primary degrdn. of the herbicide, which obeys a pseudo-first order law, was obsd. Complete mineralization of the org. C to CO2 was obtained after long-term irradiation, with corresponding stoichiometric transformation of org. Cl into Cl- ion. Various arom. intermediates, originating from 2,3,6-TBA, were detected during treatment and identified using gas chromatog.-mass spectrometry. From the anal. data, a possible multi-step degrdn. scheme is proposed. Photocatalytic treatment of the pesticide was also performed in the presence of Brij 35 micellar solns.; strong inhibition of the process was obsd. When surfactant aggregates are present, photocatalytic destruction of 2,3,6-TBA is still possible at reasonable rates only after a proper diln. of the waste and by significantly increasing the semiconductor:pollutant ratio.

IT 120-82-1, 1,2,5-Trichlorobenzene  
(anal. monitoring of surfactant effect on photocatalytic degrdn. of 2,3,6-trichlorobenzoic acid in **aq. titania dispersions**)

RN 120-82-1 HCAPLUS


CN Benzene, 1,2,4-trichloro- (8CI, 9CI) (CA INDEX NAME)



IT 9002-92-0, Brij 35  
(anal. monitoring of surfactant effect on photocatalytic degrdn. of 2,3,6-trichlorobenzoic acid in **aq. titania dispersions**)

RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



- CC 60-2 (Waste Treatment and Disposal)  
Section cross-reference(s): 5, 61, 67
- ST photocatalytic degrdn trichlorobenzoic acid **aq** titania  
**dispersion**; irradiated semiconductor photocatalysis  
pesticide water purifn; wastewater treatment irradiated  
semiconductor photocatalysis pesticide
- IT Photolysis kinetics  
(anal. monitoring of surfactant effect on photocatalytic degrdn.  
of 2,3,6-trichlorobenzoic acid in **aq**. titania  
**dispersions**)
- IT Chlorides, processes  
(anal. monitoring of surfactant effect on photocatalytic degrdn.  
of 2,3,6-trichlorobenzoic acid in **aq**. titania  
**dispersions**)
- IT Wastewater treatment  
Water purification  
(photolytic, titania-catalyzed; anal. monitoring of surfactant  
effect on photocatalytic degrdn. of 2,3,6-trichlorobenzoic acid  
in **aq**. titania **dispersions**)
- IT Solar UV radiation  
(simulated; anal. monitoring of surfactant effect on  
photocatalytic degrdn. of 2,3,6-trichlorobenzoic acid in  
**aq**. titania **dispersions**)
- IT 13463-67-7, Titania, uses  
(anal. monitoring of surfactant effect on photocatalytic degrdn.  
of 2,3,6-trichlorobenzoic acid in **aq**. titania  
**dispersions**)
- IT 95-95-4, 2,4,5-Trichlorophenol 120-82-1,  
1,2,5-Trichlorobenzene 124-38-9, Carbon dioxide, processes  
583-78-8, 2,5-Dichlorophenol 608-94-6, Trichlorohydroquinone  
933-75-5, 2,3,6-Trichlorophenol 933-78-8, 2,3,5-Trichlorophenol  
25167-80-0, Chlorophenol 25321-22-6, Dichlorobenzene  
(anal. monitoring of surfactant effect on photocatalytic degrdn.  
of 2,3,6-trichlorobenzoic acid in **aq**. titania  
**dispersions**)
- IT 9002-92-0, Brij 35  
(anal. monitoring of surfactant effect on photocatalytic degrdn.  
of 2,3,6-trichlorobenzoic acid in **aq**. titania  
**dispersions**)
- IT 50-31-7, 2,3,6-Trichlorobenzoic acid  
(anal. monitoring of surfactant effect on photocatalytic degrdn.  
of 2,3,6-trichlorobenzoic acid in **aq**. titania  
**dispersions**)

IT 7440-44-0, Carbon, processes  
(total org.; anal. monitoring of surfactant effect on  
photocatalytic degrdn. of 2,3,6-trichlorobenzoic acid in  
aq. titania **dispersions**)

L70 ANSWER 6 OF 18 HCAPLUS COPYRIGHT 2002 ACS

1998:538707 Document No. 129:293792 **Emulsion**


**polymerization** of .gamma.-benzyl L-glutamate NCA intended  
for preparation microspheres of poly(amino acid)s. Goto, Kohei;  
Yamakawa, Yoshitaka; Yoshida, Yoshinori; Hayashi, Toshio (Tsukuba  
Res. Lab., JSR Corporation, Tsukuba, 305-0841, Japan). Seitai  
Zairyo, 16(3), 145-151 (Japanese) 1998. CODEN: SEZAEH. ISSN:  
0910-304X. Publisher: Nippon Baiomateriaru Gakkai.

AB Polymn. of amino acid N-carboxy anhydrides (NCA) is usually carried  
out in anhyd. inert org. solvent, due to the tendency of amino acid  
NCA to be hydrolyzed with water. In this study we investigated the  
**emulsion polymn.** of .gamma.-benzyl-L-glutamate  
(BLG) in a mixt. of org. solvent/water in order to prep.  
microspheres of poly(amino acid). A high mol. wt.  
poly(.gamma.-benzyl-L-glutamate) (PBLG) could successfully be  
obtained with high yield by **emulsion polymn.**  
**stabilized** as O/W emulsion, when the emulsifier was a  
nonionic surfactant with high HLB and a chlorinated hydrocarbon was  
used as org. solvent, that exhibits a low soly. for water and  
appropriate dipole moment. Both factors lead to an increase of the  
polymn. rate. We also obtained the microspheres of PBLG with diam.  
of 200.apprx.300nm by distg. the org. solvent out of the emulsion,  
which is a novel and facile prepn. method of microspheres of  
poly(amino acid) derivs.

IT 9002-92-0, Polyoxyethylene lauryl ether  
(**emulsion polymn.** of .gamma.-benzyl  
L-glutamate NCA intended for prepn. microspheres of poly(amino  
acid))

RN 9002-92-0 HCAPLUS


CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



IT 67-66-3, Chloroform, processes  
(**emulsion polymn.** of .gamma.-benzyl  
L-glutamate NCA intended for prepn. microspheres of poly(amino  
acid))

RN 67-66-3 HCAPLUS

CN Methane, trichloro- (9CI) (CA INDEX NAME)



- CC 63-5 (Pharmaceuticals)  
Section cross-reference(s): 34, 35
- ST benzyl glutamate NCA **polymn emulsion** microsphere
- IT Surfactants  
(**emulsion polymn.** of .gamma.-benzyl  
L-glutamate NCA intended for prepn. microspheres of poly(amino  
acid))
- IT **Polymerization**  
(**emulsion; emulsion polymn.** of  
.gamma.-benzyl L-glutamate NCA intended for prepn. microspheres  
of poly(amino acid))
- IT Drug delivery systems  
(microspheres; **emulsion polymn.** of  
.gamma.-benzyl L-glutamate NCA intended for prepn. microspheres  
of poly(amino acid))
- IT 50-00-0D, Formaldehyde, condensate with sodium naphthalenesulfonate,  
uses 112-00-5, Lauryltrimethylammonium chloride 1338-41-6,  
Sorbitan monostearate **9002-92-0**, Polyoxyethylene lauryl  
ether 9004-96-0, Polyoxyethylene monooleate 9005-64-5,  
Polyoxyethylene sorbitan monolaurate 9005-67-8, Polyoxyethylene  
sorbitan monostearate 9016-45-9, Polyoxyethylene nonylphenyl ether  
9063-89-2, Polyoxyethylene octylphenyl ether 25155-30-0, Sodium  
dodecylbenzenesulfonate 25496-72-4, Glycerol monooleate  
37340-69-5D, Polyoxyethylene sulfate, alkyl ethers, sodium salt  
(**emulsion polymn.** of .gamma.-benzyl  
L-glutamate NCA intended for prepn. microspheres of poly(amino  
acid))
- IT 67-66-3, Chloroform, processes 75-09-2, Dichloromethane,  
processes 95-50-1, o-Dichlorobenzene 107-06-2,  
1,2-Dichloroethane, processes 108-90-7, Chlorobenzene, processes  
109-99-9, Thf, processes 141-78-6, Ethyl acetate, processes  
355-42-0, Perfluorohexane  
(**emulsion polymn.** of .gamma.-benzyl  
L-glutamate NCA intended for prepn. microspheres of poly(amino  
acid))
- IT 3190-71-4, .gamma.-Benzyl L-glutamate N-carboxyanhydride  
(**emulsion polymn.** of .gamma.-benzyl  
L-glutamate NCA intended for prepn. microspheres of poly(amino  
acid))
- IT 25014-27-1P, Poly(.gamma.-benzyl L-glutamate) 25038-53-3P  
(**emulsion polymn.** of .gamma.-benzyl  
L-glutamate NCA intended for prepn. microspheres of poly(amino  
acid))


copolymers involving steam-stripping method. Oshino, Yasuhiro; Ohtsuka, Yoshihiro (Daicel Chemical Industries, Ltd., Japan; Oshino, Yasuhiro; Ohtsuka, Yoshihiro). PCT Int. Appl. WO 9730095 A1 19970821, 39 pp. DESIGNATED STATES: W: CN, KR, US; RW: DE, ES, FR, GB, IT, NL. (Japanese). CODEN: PIXXD2. APPLICATION: WO 1996-JP334 19960216.

AB A process for producing an epoxidized block copolymer appropriate for modifying agents or modifying aids for rubbery polymers or resinous polymers, adhesives, sealants, etc. comprises epoxidizing a block copolymer which may be partly hydrated and comprises a polymer block (A) contg. a vinyl arom. hydrocarbon compd. as the main component and another polymer block (B) contg. a conjugated diene compd. as the main component to thereby synthesize an epoxidized block copolymer (C) and then recovering this polymer from a slurry or soln. thereof in an org. solvent by steam-stripping in the presence of a surfactant. It is preferably that the surfactant is a nonionic one, in particular, one represented by the general formula:  $\text{HO}(\text{C}_2\text{H}_4\text{O})_a(\text{C}_3\text{H}_6\text{O})_b(\text{C}_2\text{H}_4)_c\text{OH}$  ( $a, b, c = 1-1000$ ), having an av. mol. wt. of 3,000-20,000; and contg. 20-90 wt.% of polyoxyethylene. According to the process, the epoxidized block copolymer product can be easily recovered from the reaction mixt. without giving any prolonged heat history and thus the epoxy group can be prevented from undergoing ring opening. Also, it is possible to obtain an epoxidized block copolymer having such an epoxy equiv. as to give a high elasticity and the characteristics of epoxy compds.

IT 67-66-3, Chloroform, uses  
(solvent; manuf. of epoxidized block copolymers by steam-stripping method)

RN 67-66-3 HCAPLUS


CN Methane, trichloro- (9CI) (CA INDEX NAME)



IT 9002-92-0, Emulgen 147  
(surfactant; manuf. of epoxidized block copolymers by steam-stripping method)


RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



IC ICM C08F008-08


- CC 37-3 (Plastics Manufacture and Processing)
- IT 67-66-3, Chloroform, uses 108-88-3, Toluene, uses 110-54-3, Hexane, uses 110-82-7, Cyclohexane, uses 141-78-6, Ethyl acetate, uses 1330-20-7, Xylene, uses (solvent; manuf. of epoxidized block copolymers by steam-stripping method)
- IT 112-00-5, Quartamin 24P 7664-38-2D, Phosphoric acid, alkyl esters, compds. with alkanolamine, uses 9002-92-0, Emulgen 147 9003-11-6, Emulgen PP 290 9004-99-3, Emanon 3199 9016-45-9, Emulgen 985 26266-57-9, Rheodol SP P10 88984-51-4, Electrostripper F (surfactant; manuf. of epoxidized block copolymers by steam-stripping method)
- L70 ANSWER 8 OF 18 HCAPLUS COPYRIGHT 2002 ACS
- 1997:414049 Document No. 127:35923 Brushed fabric and textile finish. Whitley, David Anderson; Wolhar, Carl Lewis (Ivax Industries, Inc., USA). PCT Int. Appl. WO 9716259 A1 19970509, 19 pp. DESIGNATED STATES: W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, UZ, VN; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2. APPLICATION: WO 1996-US16976 19961023. PRIORITY: US 1995-550742 19951031.
- AB Natural or synthetic pile or brushed fabrics are treated to reduce marking caused by rubbing or handling the fabrics, leaving the fabrics with a soft feel and a full, thick hand. The liq. finishing compns. comprise (a) .apprx.80-100% **water-dispersible**, hydrophilic polyester or polyurethane **dispersed** in an **aq.** medium; (b) .apprx.0-20% carrier used to enhance the penetration of dyes and other materials into the polymer structure of polyester fibers; and (c) .apprx.0-10% nonrewetting wetting agent. A treatment compn. of Milease T 93, 85/15 Bu benzoate/emulsifier 4.7, and ethoxylated decyl alc. (20% soln.) 2.3% was applied to woven polyester velour upholstery fabric.
- IT 12002-48-1, Trichlorobenzene 26183-52-8, Polyethylene glycol decyl ether (in finish for brushed fabric to reduce marking caused by handling)
- RN 12002-48-1 HCAPLUS
- CN Benzene, trichloro- (8CI, 9CI) (CA INDEX NAME)



3 ( D1-C1 )

RN 26183-52-8 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-decyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IC ICM B05D003-02

ICS B32B003-02; B32B007-00; C08G018-28; C08G018-70; C08L067-02

CC 40-9 (Textiles and Fibers)

IT Polyesters, uses

Polyurethanes, uses

(water-dispersible, hydrophilic; in finish

for brushed fabric to reduce marking caused by handling)

IT 65-85-0, Benzoic acid, uses 69-72-7, Salicylic acid, uses  
 90-43-7, o-Phenyl phenol 92-52-4, Biphenyl, uses 92-69-3,  
 [1,1'-Biphenyl]-4-ol 95-57-8, o-Chlorophenol 98-85-1,  
 Methylphenyl carbinol 98-86-2, Acetophenone, uses 100-52-7,  
 Benzaldehyde, uses 108-90-7, uses 108-93-0, Cyclohexanol, uses  
 119-36-8, Methyl salicylate 119-64-2, Tetrahydronaphthalene  
 120-83-2, 2,4-Dichlorophenol 131-17-9, Diallyl phthalate  
 135-19-3, .beta.-Naphthol, uses 136-60-7, Butyl benzoate  
 513-08-6, Tripropylphosphate 567-47-5, .beta.-Naphtholsulfonic  
 acid 1321-94-4, Monomethylnaphthalene 1322-19-6  
 12002-48-1, Trichlorobenzene 25321-22-6, Dichlorobenzene  
 26183-52-8, Polyethylene glycol decyl ether 35884-90-3  
 73506-88-4 82852-79-7, Milease T

(in finish for brushed fabric to reduce marking caused by  
 handling)

L70 ANSWER 9 OF 18 HCAPLUS COPYRIGHT 2002 ACS

1996:544101 Document No. 125:177462 Surface-modified nanoparticles and  
 method of making and using them. Levy, Robert J.; Labhasetwar,  
 Vinod; Song, Cunxian S. (USA). PCT Int. Appl. WO 9620698 A2  
 19960711, 170 pp. DESIGNATED STATES: W: AL, AM, AT, AU, CA, CH,  
 CN, CZ, DE, DK, GB, HU, IS, JP, KE, LU, VN, MN, NO, US; RW: AT, BE,  
 CH, DE, ES, FR, GB, IT, LU, MR, NE, NL, PT, SE, NL, SN. (English).  
 CODEN: PIXXD2. APPLICATION: WO 1996-US476 19960104. PRIORITY: US


1995-369541 19950105; US 1995-389893 19950216.

AB Biodegradable controlled-release nanoparticles as sustained release bioactive agent delivery vehicles include surface modifying agents to target binding of the nanoparticles to tissues or cells of living systems, to enhance nanoparticle sustained release properties, and to protect nanoparticle-incorporated bioactive agents. Unique methods of making small (10 nm to 15 nm, and preferably 20 nm to 35 nm) nanoparticles having a narrow size distribution which can be surface-modified after the nanoparticles are formed is described. Techniques for modifying the surface include a lyophilization technique to produce a phys. adsorbed coating and epoxy-derivatization to functionalize the surface of the nanoparticles to covalently bind mols. of interest. The nanoparticles may also comprise hydroxy-terminated or epoxide-terminated and/or activated multiblock copolymers, having hydrophobic segments which may be polycaprolactone and hydrophilic segments. The nanoparticles are useful for local intravascular administration of smooth muscle inhibitors and antithrombogenic agents as part of interventional cardiac or vascular catheterization such as a balloon angioplasty procedure; direct application to tissues and/or cells for gene therapy, such as the delivery of osteotropic genes or gene segments into bone progenitor cells; or oral administration in an enteric capsule for delivery of protein/peptide based vaccines.

IT 67-66-3, Chloroform, biological studies  
(solvent; surface-modified polymer controlled-release nanoparticles for sustained drug delivery)

RN 67-66-3 HCAPLUS


CN Methane, trichloro- (9CI) (CA INDEX NAME)



IT 75-47-8, Iodoform  
(surface-modified polymer controlled-release nanoparticles for sustained drug delivery)

RN 75-47-8 HCAPLUS

CN Methane, triiodo- (8CI, 9CI) (CA INDEX NAME)




IT 9002-92-0, Polyoxyethylene lauryl ether  
(surface-modified polymer controlled-release nanoparticles for sustained drug delivery)

RN 9002-92-0 HCAPLUS



CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



IC A61K009-51

CC 63-6 (Pharmaceuticals)

IT 67-64-1, 2-Propanone, biological studies 67-66-3,  
Chloroform, biological studies 67-68-5, Dimethylsulfoxide,  
biological studies 68-12-2, Dimethylformamide, biological studies  
75-09-2, Methylene chloride, biological studies 109-99-9,  
biological studies 123-91-1, Dioxane, biological studies  
127-19-5, Dimethylacetamide 141-78-6, Ethyl acetate, biological  
studies 684-16-2, Hexafluoroacetone 920-66-1

(solvent; surface-modified polymer controlled-release  
nanoparticles for sustained drug delivery)

IT 75-23-0 75-47-8, Iodoform 102-54-5, Ferrocene  
113-00-8, Guanidine 288-32-4, Imidazole, uses 558-13-4, Carbon  
tetrabromide 7550-45-0, Titanium tetrachloride, uses 7637-07-2D,  
Boron trifluoride, adducts 13598-36-2D, Phosphonic acid,  
alkylidenebis- derivs. 13826-88-5, Zinc tetrafluoroborate  
86665-14-7, Zirconocene chloride

(surface-modified polymer controlled-release nanoparticles for  
sustained drug delivery)

IT 50-70-4, D-Glucitol, biological studies 57-09-0, Cetyl trimethyl  
ammonium bromide 57-10-3, Hexadecanoic acid, biological studies  
57-88-5, Cholesterol, biological studies 69-65-8, D-Mannitol  
102-71-6, Triethanolamine, biological studies 112-02-7, Hexadecyl  
trimethyl ammonium chloride 151-21-3, Sodium dodecyl sulfate,  
biological studies 577-11-7, Sodium dioctyl sulfosuccinate  
1069-55-2, Isobutyl cyanoacrylate 3282-73-3, Didodecyldimethyl  
ammonium bromide 7445-62-7 7727-43-7, Barium sulfate  
8007-43-0, Sorbitan sesquioleate 9000-65-1, Tragacanth  
9000-69-5, Pectin 9002-89-5, Polyvinyl alcohol 9002-92-0  
, Polyoxyethylene lauryl ether 9003-39-8, Polyvinyl pyrrolidone  
9003-53-6, Polystyrene 9004-32-4 9004-34-6, Cellulose,  
biological studies 9004-35-7, Cellulose acetate 9004-44-8,  
Cellulose phthalate 9004-64-2, Hydroxypropyl cellulose 9004-99-3  
9005-49-6, Heparin, biological studies 9015-73-0 9050-04-8,  
CM-cellulose calcium 9050-31-1, Hydroxypropyl methyl cellulose  
phthalate 10103-46-5, Calcium phosphate 25322-68-3  
106392-12-5, Poloxamer 110617-70-4, Poloxamine 128835-92-7,  
Lipofectin 180741-27-9

(surface-modified polymer controlled-release nanoparticles for  
sustained drug delivery)

IT 7732-18-5, **Water**, biological studies  
(**suspending** medium; surface-modified polymer  
controlled-release nanoparticles for sustained drug delivery)

L70 ANSWER 10 OF 18 HCAPLUS COPYRIGHT 2002 ACS

1994:540284 Document No. 121:140284 Coated substrates and laminate structures comprising organic solvent-based dispersions of organocation-modified vermiculite.. Ou, Chia Chih; Bablouzian, Leon (W.R. Grace and Co.-Conn., USA). Eur. Pat. Appl. EP 601877 A1 19940615, 10 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE. (English). CODEN: EPXXDW. APPLICATION: EP 1993-309981 19931210. PRIORITY: US 1992-989262 19921211.


AB Coated substrates, and laminate structures consisting of intermediate layers provided by org. solvent-based dispersions of org. cation-modified delaminated vermiculite are disclosed. The laminates display better peel strength compared to laminates made from **water-based dispersions**, and can be used, e.g., for fire-resistant coatings, sealing gaskets, and gas barriers.

IT 67-66-3, Chloroform, uses 60828-78-6, Tergitol TMN-6

(in manuf. of vermiculite laminate building materials)


RN 67-66-3 HCAPLUS

CN Methane, trichloro- (9CI) (CA INDEX NAME)



RN 60828-78-6 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-[3,5-dimethyl-1-(2-methylpropyl)hexyl]-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IC ICM C04B014-20

ICS C08K003-34; C08K007-26

CC 58-4 (Cement, Concrete, and Related Building Materials)

IT 57-09-0, Cetyltrimethylammonium bromide 67-56-1, Methanol, uses 67-66-3, Chloroform, uses 78-93-3, Methyl ethyl ketone, uses 98-95-3, Nitrobenzene, uses 100-52-7, Benzaldehyde, uses 106-42-3, p-Xylene, uses 108-88-3, Toluene, uses 109-99-9, Tetrahydrofuran, uses 111-87-5, Octanol, uses 127-19-5, Dimethylacetamide 141-78-6, Ethyl acetate, uses 1318-00-9, Vermiculite 1330-20-7, Xylene, uses 24937-78-8, Ethylene-vinyl acetate copolymer 25038-32-8, Isoprene-styrene copolymer 39405-74-8, Terphane 60828-78-6, Tergitol TMN-6

(in manuf. of vermiculite laminate building materials)

L70 ANSWER 11 OF 18 HCAPLUS COPYRIGHT 2002 ACS

1991:512798 Document No. 115:112798 Acceleration of yeast autolysis by chemical methods for production of intracellular enzymes. Breddam, Klaus; Beenfeldt, Thorkild (Dep. Chem., Carlsberg Lab., Copenhagen Valby, DK-2500, Den.). Applied Microbiology and Biotechnology, 35(3), 323-9 (English) 1991. CODEN: AMBIDG. ISSN: 0175-7598.


AB Known methods for the acceleration of yeast autolysis were investigated and new methods were developed. Autolysis was induced by plasmolysis with a no. of solvents. The efficiency of this treatment is dependent on the nature of the solvent, its concn., and the duration of the treatment. Plasmolysis generally does not cause release of mols. of high mol. wt. (MW), such as enzymes. However, addn. of water initiates autolysis and the enzyme carboxypeptidase Y (MW 64,000), for example, is released. The rate of this process is very dependent on pH; at the optimal pH (.apprx.8.0) essentially complete autolysis is achieved within 20 h using the best solvents. Control of pH through the process is required. Straight-chain alcs. of medium chain length, i.e. C6-C9, appear to function efficiently in amts. of only 1.2 mL/100 g yeast. In amts. of 2.5-10 mL solvent/100 g yeast, trichloroethane, CHCl<sub>3</sub>, and, in particular, ether also provide efficient plasmolysis. Treatment of an **aq. suspension** of yeast cells with a variety of nonionic as well as ionic detergents caused autolysis. The effect of pH corresponds to that obsd. with org. solvents, i.e., a pH of .apprx.8.0 is optimal. This autolysis was most efficient when the compressed yeast had been initially plasmolyzed by treatment with NaCl followed by addn. of water. The inexpensive detergents Triton X-100 and N-lauroylsarcosine appeared to be among the most efficient. These methods are inexpensive and can be employed on a large scale. In addn., cell debris is easily removed, which is very important for subsequent downstream processing. In the alternative method using phys. breakage by homogenization this step is highly problematic.

IT 25323-89-1, Trichloroethane 60828-78-6, Tergitol  
TMN6

(yeast autolysis by, for prodn. of intracellular enzymes)

RN 25323-89-1 HCAPLUS

CN Ethane, trichloro- (8CI, 9CI) (CA INDEX NAME)




D1-C1


RN 60828-78-6 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-[3,5-dimethyl-1-(2-

methylpropyl)hexyl]-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IT 67-66-3, Chloroform, reactions  
 (yeast autolysis by, for prodn. of intracellular enzymes)  
 RN 67-66-3 HCAPLUS  
 CN Methane, trichloro- (9CI) (CA INDEX NAME)




CC 16-9 (Fermentation and Bioindustrial Chemistry)  
 IT 25323-89-1, Trichloroethane 60828-78-6, Tergitol  
 TMN6 75621-03-3  
 (yeast autolysis by, for prodn. of intracellular enzymes)  
 IT 56-23-5, Carbon tetrachloride, reactions 57-09-0,  
 Hexadecyltrimethyl ammonium bromide 67-66-3, Chloroform,  
 reactions 71-41-0, 1-Pentanol, reactions 75-09-2,  
 Dichloromethane, reactions 81-24-3, Taurocholic acid 97-78-9  
 111-13-7, 2-Octanone 111-27-3, 1-Hexanol, reactions 111-70-6,  
 1-Heptanol 111-87-5, 1-Octanol, reactions 112-30-1, 1-Decanol  
 112-53-8, 1-Dodecanol 123-03-5, Cetylpyridinium chloride  
 143-08-8, 1-Nonanol 591-78-6, 2-Hexanone 593-51-1D, trialkyl  
 derivs. 9002-93-1, Triton X 101 9016-45-9, Tergitol NP40  
 9036-19-5  
 (yeast autolysis by, for prodn. of intracellular enzymes)


L70 ANSWER 12 OF 18 HCAPLUS COPYRIGHT 2002 ACS  
 1990:520725 Document No. 113:120725 Indomethacin polymeric  
 nanosuspensions prepared by microfluidization. Bodmeier, Roland;  
 Chen, Huagang (Coll. Pharm., Univ. Texas, Austin, TX, 78712-1074,  
 USA). Journal of Controlled Release, 12(3), 223-33 (English) 1990.  
 CODEN: JCREEC. ISSN: 0168-3659.  
 AB Polymeric nanosuspensions contg. indomethacin were prepd. by a  
 microfluidization-solvent evapn. method. The nanosuspensions were  
 evaluated with respect to total drug content, drug content in the  
 polymer and aq. phase, particle size, drug crystn. in the aq. phase,  
 in vitro drug release, and stability to flocculation in 0.1N HCl and  
 pH 7.4 phosphate buffer. Nanosuspensions with a total drug content  
 of 35 mg indomethacin/mL nanosuspension could be prepd. without drug  
 crystn. More than 98.permill. of the drug were found within the  
 polymer phase. Unwanted drug crystn. in the aq. phase depended on  
 the drug loading, the drug-polymer compatibility, the org. solvent,

and the type and amt. of surfactant used. Indomethacin was released from Et cellulose nanoparticles within 15 min. Nanoparticles intended to provide drug release over longer periods of time were obtained by using mixts. of Et cellulose and poly(Me methacrylate). In pH 7.4 buffer, anionic, nonionic, and macromol. stabilizers protected the nanosuspensions against flocculation while nonionic surfactants were good stabilizers in 0.1N HCl.


IT 9002-92-0 9005-00-9, Brij 78  
(Et cellulose nanosuspension contg., indomethacin stability in relation to)  
RN 9002-92-0 HCAPLUS  
CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



RN 9005-00-9 HCAPLUS  
CN Poly(oxy-1,2-ethanediyl), .alpha.-octadecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



IT 67-66-3, uses and miscellaneous  
(indomethacin dissoln. from nanosuspension in relation to)  
RN 67-66-3 HCAPLUS  
CN Methane, trichloro- (9CI) (CA INDEX NAME)



CC 63-6 (Pharmaceuticals)  
IT Particle size  
(of **polymer** nanoparticles for **suspension**,  
sodium lauryl sulfate and pressure effect on)  
IT 9002-89-5, Poly(vinyl alcohol) 9002-92-0 9004-99-3  
9005-00-9, Brij 78 9005-65-6, Tween 80 106392-12-5  
(Et cellulose nanosuspension contg., indomethacin stability in relation to)  
IT 67-66-3, uses and miscellaneous 75-09-2, uses and  
miscellaneous 141-78-6, Acetic acid ethyl ester, uses and  
miscellaneous

(indomethacin dissoln. from nanosuspension in relation to)

L70 ANSWER 13 OF 18 HCAPLUS COPYRIGHT 2002 ACS


1987:460254 Document No. 107:60254 Manufacture of flame-retardant foams from urethane **polymer emulsions**. Mai, Kazumi; Midorikawa, Akio; Takegawa, Hisao; Kawanami, Eiji (Dainippon Ink and Chemicals, Inc., Japan). Jpn. Kokai Tokkyo Koho JP 62041237 A2 19870223 Showa, 11 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 1985-179462 19850816.

AB Title foams having good abrasion and light resistance, useful esp. for automobile seats, are manufd. by mech. foaming mixed emulsions of polyurethanes, conjugated diene and/or vinyl copolymers, ethylene oxide and/or propylene oxide polymers, and flame retardants, then spreading on substrates or casting in molds, and heating. Thus, a 1,6-hexamethylenediamine-polyoxypropylene glycol-TDI **copolymer (I) emulsion** was mixed with Tergitol XD (ethylene oxide-propylene oxide copolymer), acrylonitrile-butadiene-methacrylic acid **copolymer emulsion**, **emulsified** C6Br5OC6Br5 and Sb2O5, and other additives, mech. foamed, spread on polyester fabric, and heated at 120.degree. for 8 min to form a uniform, smooth foam sheet which passed the MVSS 302 flame retardance test, and showed good abrasion and light resistance, in contrast to one prepd. without the I and Tergitol XD.

IT **25637-99-4**, Hexabromocyclododecane  
(flame retardants, polyurethane-polyoxyalkylene-vinyl/diene copolymer blend foams contg., manufd. from mixed emulsions)

RN 25637-99-4 HCAPLUS

CN Cyclododecane, hexabromo- (7CI, 8CI, 9CI) (CA INDEX NAME)



6 ( D1- Br )

IT **9038-95-3**, Tergitol XD  
(polyurethane-vinyl/diene **copolymer** mixed **emulsions** contg, with flame-retardants, foams manufd. from)

RN 9038-95-3 HCAPLUS

CN Oxirane, methyl-, polymer with oxirane, monobutyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 71-36-3

CMF C4 H10 O



CM 2

CRN 9003-11-6


CMF (C3 H6 O . C2 H4 O)x

CCI PMS

CM 3

CRN 75-56-9

CMF C3 H6 O



CM 4

CRN 75-21-8

CMF C2 H4 O



- IC ICM C08J009-30  
 CC 38-3 (Plastics Fabrication and Uses)  
 IT Polyoxyalkylenes, uses and miscellaneous  
     (polyurethane-vinyl/diene **copolymer** mixed  
     **emulsions** contg, with flame-retardants, foams manufd.  
     from)  
 IT 1163-19-5, Decabromodiphenyl ether 1314-60-9, Antimony pentaoxide  
     6145-73-9 **25637-99-4**, Hexabromocyclododecane 36704-02-6  
     (flame retardants, polyurethane-polyoxyalkylene-vinyl/diene  
     copolymer blend foams contg., manufd. from mixed emulsions)  
 IT **9038-95-3**, Tergitol XD  
     (polyurethane-vinyl/diene **copolymer** mixed  
     **emulsions** contg, with flame-retardants, foams manufd.  
     from)


1985:134011 Document No. 102:134011 Soil-resistant detergents for carpets. (Tokyo Livin K. K., Japan). Jpn. Kokai Tokkyo Koho JP 59196400 A2 19841107 Showa, 6 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 1983-70145 19830422.

AB The detergents effective in prevention of soil redeposition for carpets surface-processed with F-compds. contain 0.01-0.4% nonionic surfactant and org. solvent (soly. .gtoreq.2% in water at 25.degree.) and water (in total 100%), and after drying solids other than the nonionic surfactant are negligible. The detergents remain in very small amts. after cleaning. Thus, a 5% **water dispersion** contg. 38% perfluoroalkyl group-contg. acrylate oligomer was spray-coated on a nylon 66 carpet (2400 denier), heated 14 min at 145.degree., and cut to prep. sample pieces (10 cm .times. 10 cm), which were washed by spraying (0.15 g/m2) with a detergent contg. poly(ethylene oxide) lauryl ether [9002-92-0] (d.p. 6) 0.05, MEK [78-93-3] 10, and water 89.95% and wiped off, showing good cleaning effect, soil resistance, and little redeposition.

IT 79-01-6, uses and miscellaneous 9002-92-0  
(cleaning compns. contg., for carpets)


RN 79-01-6 HCAPLUS

CN Ethene, trichloro- (9CI) (CA INDEX NAME)



RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



IC C11D010-02

CC 46-6 (Surface Active Agents and Detergents)

IT 71-36-3, uses and miscellaneous 78-83-1, uses and miscellaneous  
78-93-3, uses and miscellaneous 79-01-6, uses and  
miscellaneous 111-76-2 141-78-6, uses and miscellaneous  
9002-92-0 9003-11-6 9004-99-3 9016-45-9 25322-68-3D,  
ethers with sperm oil alcs.  
(cleaning compns. contg., for carpets)


L70 ANSWER 15 OF 18 HCAPLUS COPYRIGHT 2002 ACS

1975:581045 Document No. 83:181045 Composition making textiles oil- and water-repellent. Inman, Charles E. (Pennwalt Corp., USA). Ger. Offen. DE 2460142 19750710, 26 pp. (German). CODEN: GWXXBX.




APPLICATION: DE 1974-2460142 19741219.

- AB Emulsions for oil- and waterproofing textiles contain a chlorinated C1-3 alkane solvent, a small amt. of **water**, a finely **dispersed** fluoroalkyl **polymer** with C6-16 fluoroalkyl groups, and an **emulsion-stabilizing** nonionic surfactant with hydrophile-lipophile value 8-20. For example, a **water-in-oil emulsion** was prepd. by mixing methylchloroform [71-55-6] 100, Tween 40 [9005-66-7] surfactant 1, and fluoropolymer latex 25 g (contg. 20% polymer mixt. of 50 parts 80:20 C9F19CH2CH2SCOCMe:CH2-stearyl methacrylate polymer [30661-93-9] and 50 parts 71:5:24 butyl methacrylate-N-methylolacrylamide-3,5,5-trimethylhexyl methacrylate polymer [56993-04-5]). Cotton, polyester, acrylic, nylon, and cotton-polyester textiles finished with the emulsion had good oil and water repellency.
- IT **71-55-6**  
(solvents, fluoropolymer **water-in-oil emulsions** contg., for oil- and waterproofing of textiles)
- RN 71-55-6 HCAPLUS
- CN Ethane, 1,1,1-trichloro- (8CI, 9CI) (CA INDEX NAME)




- IT **9005-00-9**  
(surfactants, fluoropolymer **water-in-oil emulsions** contg., for oil- and waterproofing of textiles)
- RN 9005-00-9 HCAPLUS
- CN Poly(oxy-1,2-ethanediyl), .alpha.-octadecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)




- IC D06M
- CC 39-10 (Textiles)
- IT Oils  
(-proofing, of textiles, fluoropolymer **water-in-oil emulsions** for)
- IT Surfactants  
(nonionic polyethylene glycol derivs., fluoropolymer **water-in-oil emulsions** contg., for oil- and waterproofing of textiles)
- IT Waterproofing


- (of textiles, fluoropolymer **water-in-oil emulsions** for)
- IT Textiles  
(oil- and waterproofing of cotton, fluoropolymer **water-in-oil emulsions** for)
- IT Acrylic fibers  
Polyamide fibers  
Polyester fibers  
(oil- and waterproofing of, fluoropolymer **water-in-oil emulsions** for)
- IT Poly(oxy-1,2-ethanediyl), .alpha.-hydro-.omega.-hydroxy-, ethers, fatty acid glyceride derivs.  
(surfactants, fluoropolymer **water-in-oil emulsions** contg., for oil- and waterproofing of textiles)
- IT 71-55-6  
(solvents, fluoropolymer **water-in-oil emulsions** contg., for oil- and waterproofing of textiles)
- IT 1338-43-8 9004-98-2 9004-99-3 9005-00-9 9005-65-6  
9005-66-7 9005-67-8 31694-55-0  
(surfactants, fluoropolymer **water-in-oil emulsions** contg., for oil- and waterproofing of textiles)
- L70 ANSWER 16 OF 18 HCAPLUS COPYRIGHT 2002 ACS  
1973:443846 Document No. 79:43846 Laundry detergent composition. Marple, Walter L. (Whirlpool Corp.). U.S. US 3737387 19730605, 3 pp. (English). CODEN: USXXAM. APPLICATION: US 1970-46278 19700615.
- AB Soils contg. complex fats, e.g. lard and butter, were efficiently removed from fabrics made from natural or synthetic fibers with a detergent compn. contg. 40-80% by vol. of a **water-sol. emulsifier** and 60-20% by vol. of an org. solvent. Thus, a mixt. of 40 vol.% polyethylene glycol mono(nonylphenyl) ether [9016-45-9] and 60 vol.% trichloroethylene [79-01-6] was dild. to a 0.2% aq. soln. A fabric soiled with lard was washed 5 min in a conventional home laundering detergent, then 5 min in the above soln. to give 77.3% lard removal. Increasing the 2nd cycle to 10 min removed 87.3% of the lard, compared with 42.0% lard removal after washing for 10 min with a 0.2% soln. of a conventional detergent.
- IT 71-55-6 79-01-6  
(cleaning **emulsions** contg. **water** and, for textiles)
- RN 71-55-6 HCAPLUS  
CN Ethane, 1,1,1-trichloro- (8CI, 9CI) (CA INDEX NAME)



RN 79-01-6 HCAPLUS  
 CN Ethene, trichloro- (9CI) (CA INDEX NAME)



IT 24938-91-8  
 (emulsifying agents, for water-solvent  
 cleaning compns., for textiles)  
 RN 24938-91-8 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-tridecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



IC C09D; C11D; C23G  
 NCL 252170000  
 CC 46-6 (Surface Active Agents and Detergents)  
 ST polyethylene glycol chloroethylene detergent; lard removal soiled  
 fabric; butter removal soiled fabric; **water sol**  
**emulsifier** detergent; org solvent emulsifier detergent;  
 laundry detergent system  
 IT 71-55-6 79-01-6 108-32-7 127-18-4  
 (cleaning **emulsions** contg. **water** and, for  
 textiles)  
 IT 9003-11-6 9004-81-3 9004-87-9 9016-45-9 24938-91-8  
 (emulsifying agents, for **water**-solvent  
 cleaning compns., for textiles)  
 L70 ANSWER 17 OF 18 HCAPLUS COPYRIGHT 2002 ACS  
 1972:42770 Document No. 76:42770 Highly viscous pesticidal dispersions  
 for spraying. Battani, Noel; Geiger, Max (Schloesing S. A;  
 Ciba-Geigy A.-G.). Ger. Offen. DE 2109410 19711007, 53 pp.  
 (German). CODEN: GWXXBX. APPLICATION: DE 1971-2109410 19710227.  
 AB Pesticidal compns. were stored as oil-in-**water**  
**dispersions**, and were converted before use into highly  
 viscous **water**-in-oil **dispersions**, by addn. of a  
 dispersing agent in a polar solvent. Thus, a storable insecticidal  
 oil-in-**water dispersion** contg. **H2O**  
 230, O,O-dimethyl O-(4-nitro-m-tolyl) phosphorothioate [122-14-5]  
 74, BuOAc 25, gas oil 335, 1:4 C12-14 alc.-ethylene oxide  
 condensation product 12, 1:9 nonylphenol-ethylene oxide condensation  
 product [39587-22-9] 3, CaCO3 375, and kaolin 150 g, was  
 treated before use with 20 g fatty acid-fatty amine salt complex and


20 g propylene glycol [57-55-6] to give a 1 l. dispersion suitable for spraying.

IT 71-55-6

(dispersing agents for pesticides)

RN 71-55-6 HCAPLUS

CN Ethane, 1,1,1-trichloro- (8CI, 9CI) (CA INDEX NAME)




IT 58-89-9

(oil-in-water dispersions of, manufg. of viscous)

RN 58-89-9 HCAPLUS

CN Cyclohexane, 1,2,3,4,5,6-hexachloro-, (1.alpha.,2.alpha.,3.beta.,4.alpha.,5.alpha.,6.beta.) - (9CI) (CA INDEX NAME)

Relative stereochemistry.




IT 9002-92-0

(pesticide dispersion in oligomeric, prepn. of viscous)

RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)




IT 27044-11-7

(pesticide dispersions in, prepn. of viscous)

RN 27044-11-7 HCAPLUS


IC A01N

- CC 5 (Agrochemicals)  
 IT Pesticides  
     (oil-in-water dispersions of, manufg. of viscous)  
 IT 57-55-6, biological studies 71-55-6 108-94-1, biological studies  
     (dispersing agents for pesticides)  
 IT 58-89-9 122-14-5 137-26-8 900-95-8 1344-71-4  
     8018-01-7 12122-67-7 35565-86-7  
     (oil-in-water dispersions of, manufg. of viscous)  
 IT 9002-92-0  
     (pesticide dispersion in oligomeric, prepn. of viscous)  
 IT 9003-11-6 9056-20-6 27044-11-7  
     (pesticide dispersions in, prepn. of viscous)
- L70 ANSWER 18 OF 18 HCAPLUS COPYRIGHT 2002 ACS  
 1967:45781 Document No. 66:45781 Treating the bark of trees to prevent insect damage by tree bark beetles and insects. Stewart, Katherine Ferguson (Stewart Operations, Inc.). U.S. US 3298913 19670117, 4 pp. (English). CODEN: USXXAM. APPLICATION: US 19650901.
- AB The efficacy of insecticides such as benzene hexachloride and DDT against the tree bark beetle is enhanced by formulating a 1-5% aq. dispersion of the insecticide with 1-5% of a multicomponent surface active concentrate. The concentrate consists of a 596 mol. wt. condensate of ethylene oxide and tridecyl alc. 15-25, Na N-methyl-N-oleoyl taurate 1-2, MeOH or EtOH 2.5-3, ethylene glycol 30-5, Na2CrO4 0.05-0.5, Me, Et, or Bu Cellosolve 0.04-3, and H2O 25-45%.
- IT 50-29-3, biological studies 58-89-9, biological studies 32126-89-9  
     (insecticide compn. contg.)
- RN 50-29-3 HCAPLUS  
 CN Benzene, 1,1'-(2,2,2-trichloroethylidene)bis[4-chloro- (9CI) (CA INDEX NAME)



- RN 58-89-9 HCAPLUS  
 CN Cyclohexane, 1,2,3,4,5,6-hexachloro-, (1.alpha.,2.alpha.,3.beta.,4.alpha.,5.alpha.,6.beta.)- (9CI) (CA INDEX NAME)

Relative stereochemistry.



RN 32126-89-9 HCAPLUS  
 NCL 167042000  
 CC 19 (Pesticides)  
 IT 50-29-3, biological studies 58-89-9, biological  
 studies 110-80-5 137-20-2 7775-11-3 32126-89-9  
 (insecticide compn. contg.)


=> d 171 1-24 cbib abs hitstr hitind

L71 ANSWER 1 OF 24 HCAPLUS COPYRIGHT 2002 ACS  
 2002:539870 Document No. 137:106051 Nucleic acid extraction solution  
 and use thereof. Lentricchia, Brian; Cohenford, Menashi A. (Cytoc  
 Corporation, USA). PCT Int. Appl. WO 2002055739 A2 20020718, 30 pp.  
 DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR,  
 BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI,  
 GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,  
 LC, LK, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM,  
 PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,  
 UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM;  
 RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA,  
 GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR.  
 (English). CODEN: PIXXD2. APPLICATION: WO 2002-US1430 20020115.  
 PRIORITY: US 2001-PV261845 20010115.


AB Disclosed are methods and compns. for extg. nucleic acids from a  
 biol. sample. In particular, disclosed is a nucleic acid extn.  
 soln. together with method using such a soln. for extg. nucleic acid  
 sequences from biol. samples contg. cells, cellular debris or both.  
 The nucleic acid extn. soln. contains a mol. having the formula  
 R1O-CH2-CH2-OR2, wherein R1 and R2 independently are selected from  
 the group consisting of hydrogen and an alkyl group. Vaginal swab  
 samples spiked with Neisseria gonorrhoeae were extd. with 1 %  
 2-methoxyethanol in 2 mM borate buffer, pH 9.5.

IT 9002-92-0, Brij 35  
 (extn. soln. contg.; nucleic acid extn. soln. and use thereof)

RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



IT 67-66-3, Chloroform, miscellaneous  
 (nucleic acid extn. soln. and use thereof)  
 RN 67-66-3 HCAPLUS  
 CN Methane, trichloro- (9CI) (CA INDEX NAME)



IC ICM C12Q001-68  
 ICS C12N015-10  
 CC 9-9 (Biochemical Methods)  
 Section cross-reference(s): 1, 3, 10, 14  
 IT Biological materials  
 Cell  
 Dissolution  
 Extraction  
 Heating  
 Human  
 Human papillomavirus  
 Nucleic acid amplification (method)  
 PCR (polymerase chain **reaction**)  
 pH  
 (nucleic acid extn. soln. and use thereof)  
 IT 77-86-1, Tris buffer 107-21-1, Ethylene glycol, uses 109-86-4,  
 2-Methoxyethanol 110-80-5, 2-Ethoxyethanol 111-76-2 589-35-5  
 593-84-0, Guanidinium thiocyanate 624-95-3 1132-61-2, MOPS  
 1333-73-9, Sodium borate 4439-24-1 **9002-92-0**, Brij 35  
 9002-93-1, Triton X-100 9005-64-5, Tween 20 16484-86-9,  
 1,2-Diethoxyethene  
 (extn. soln. contg.; nucleic acid extn. soln. and use thereof)  
 IT 67-66-3, Chloroform, miscellaneous 108-95-2, Phenol,  
 miscellaneous  
 (nucleic acid extn. soln. and use thereof)

L71 ANSWER 2 OF 24 HCAPLUS COPYRIGHT 2002 ACS  
 2002:487712 Document No. 137:64931 Branched **reaction**  
 products of amines and multifunctional compounds for aqueous  
 compositions. Gross, Stephen F.; Li, Wei; Mao, Jianhua; Tuller,  
 Norman; Wiggins, Michael S. (Cognis Corporation, USA). PCT Int.  
 Appl. WO 2002050235 A1 20020627, 13 pp. DESIGNATED STATES: W: AE,  
 AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR,  
 CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
 ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,

MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2.


APPLICATION: WO 2001-US44739 20011129. PRIORITY: US 2000-PV256375 20001218; US 2001-PV281472 20010404; US 2001-975357 20011011.

AB **Reaction** products of (A) an amine  $RX(EO)_n(PO)_m(BO)_pZNHR_1$  where R = substituted or unsubstituted, satd. or unsatd., org. group having 4-36 C atoms,  $R_1 = H$  or a C1-20 straight or branched chain alkyl, optionally substituted with OH group, X = O, S, or  $NR_3$  where  $R_3 = H$  or a C1-4-alkyl, Z = ethylene, propylene or butylene, n = 0-100, m = 0-50, and p = 0-50, provided  $n+m+p$  is  $\geq 1$ ; and (B) a multifunctional compd.  $R_2(Y)_m$ , where  $R_2$  = straight or branched chain alkyl, alkenyl, or cycloaliph. group having 2-30 C atoms, or an unsubstituted or alkyl substituted arom. group, Y = carboxyl, carboxylic anhydride, halogen, carbonyl, acetyl halide, keto, aldehyde, or epoxy, and m = 2-10, and Y can be the same or different; give defoaming agents for liq. compns. (aq. and nonaq.).

IT **79-00-5DP**, 1,1,2-Trichloroethane, **reaction** products with polyethylene glycol dodecyl ether and propylamine **9002-92-0DP**, Polyethylene glycol dodecyl ether, **reaction** products with propylamine and trichloroethane (branched **reaction** products of amines and multifunctional compds. as low foaming surfactants or defoaming agents for liqs.)


RN 79-00-5 HCAPLUS

CN Ethane, 1,1,2-trichloro- (8CI, 9CI) (CA INDEX NAME)



RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IC ICM C11D017-00

ICS C11D001-825

CC 46-3 (Surface Active Agents and Detergents)

IT Antifoaming agents

(branched **reaction** products of amines and multifunctional compds. as low foaming surfactants or defoaming agents for liqs.)



## IT Surfactants

(nonionic, low foaming; branched **reaction** products of amines and multifunctional compds. as low foaming surfactants or defoaming agents for liqs.)

- IT 79-00-5DP, 1,1,2-Trichloroethane, **reaction** products with polyethylene glycol dodecyl ether and propylamine 107-10-8DP, Propylamine, **reaction** products with trichloroethane and polyethylene glycol dodecyl ether 2404-44-6DP, 1,2-Epoxydecane, **reaction** products with polyethylene glycol decyl ether oxypropylamine and ethylenediaminetetraacetic dianhydride 9002-92-0DP, Polyethylene glycol dodecyl ether, **reaction** products with propylamine and trichloroethane 23911-25-3DP, Ethylenediaminetetraacetic dianhydride, **reaction** products with polyethylene glycol decyl ether oxypropylamine and epoxydecane 439152-57-5DP, **reaction** products with epoxydecane and ethylenediaminetetraacetic dianhydride (branched **reaction** products of amines and multifunctional compds. as low foaming surfactants or defoaming agents for liqs.)

L71 ANSWER 3 OF 24 HCAPLUS COPYRIGHT 2002 ACS


2002:412950 Document No. 137:267631 The mechanisms of rate enhancing and quenching of trichloroethene photodecay in the presence of sensitizer and hydrogen sources. Chu, W.; Choy, W. K. (Research Centre for Urban Environmental Technology and Management, Department of Civil and Structural Engineering, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong). Water Research, 36(10), 2525-2532 (English) 2002. CODEN: WATRAG. ISSN: 0043-1354. Publisher: Elsevier Science Ltd..

AB The **reaction** mechanisms and rates of trichloroethene (TCE) photodecay in the presence of photosensitizer (acetone, ACE) and hydrogen sources (surfactant and triethylamine, TEA) were investigated. Quantum yields of TCE photodecay in soln. with surfactant Brij 35 and optimal ACE dosage are about 25 times higher than in Brij 35 alone. However, with an excess ACE dosage, ACE will act as a light barrier and attenuate the light intensity available for TCE photodegrdn. TCE photodegrdn. follows a two-stage kinetics, in which a lag-phase is followed by a fast decay. The lag-phase distribution depends on initial pH levels and ACE concns. The overall TCE removal was found to be higher at high pH level, suggesting that free radical **reaction** is dominant at high pH levels. The use of addnl. hydrogen source (TEA) in the **reaction** can further accelerate the **reaction**, but overdosing of TEA would quench the **reaction**. The possible **reaction** mechanisms of TCE photodecay involving ACE and TEA were proposed, and rate-enhancing and rate-quenching models at low and high TEA concns. resp. were derived based on the proposed mechanism; they were found useful for predicting the TEC decay quantum yields.


## IT 79-01-6, Trichloroethene, processes

(mechanisms of rate enhancing and quenching of trichloroethene

photodecay in soln. with sensitizer and hydrogen sources)  
 RN 79-01-6 HCAPLUS  
 CN Ethene, trichloro- (9CI) (CA INDEX NAME)




IT 9002-92-0, Brij 35  
 (surfactant, hydrogen source; mechanisms of rate enhancing and quenching of trichloroethene photodecay in soln. with sensitizer and hydrogen sources)  
 RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




- CC 60-2 (Waste Treatment and Disposal)  
 Section cross-reference(s): 19, 74
- IT 79-01-6, Trichloroethene, processes  
 (mechanisms of rate enhancing and quenching of trichloroethene photodecay in soln. with sensitizer and hydrogen sources)
- IT 67-64-1, Acetone, reactions  
 (photosensitizer; mechanisms of rate enhancing and quenching of trichloroethene photodecay in soln. with sensitizer and hydrogen sources)
- IT 121-44-8, Triethylamine, reactions 9002-92-0,  
 Brij 35  
 (surfactant, hydrogen source; mechanisms of rate enhancing and quenching of trichloroethene photodecay in soln. with sensitizer and hydrogen sources)
- L71 ANSWER 4 OF 24 HCAPLUS COPYRIGHT 2002 ACS  
 2001:920927 Document No. 136:267783 The study of rate improvement of trichloroethene (TCE) decay in UV system with hydrogen source. Choy, W. K.; Chu, W. (Research Centre for Urban Environmental Technology and Management, Department of Civil and Structural Engineering, The Hong Kong Polytechnic University, Hong Kong, Hong Kong). Water Science and Technology, 44(6, Managing Water and Waste in the New Millennium), 27-33 (English) 2001. CODEN: WSTED4. ISSN: 0273-1223. Publisher: IWA Publishing.
- AB The photosensitization of trichloroethene (TCE) in the presence of hydrogen source of surfactant and photosensitizer was investigated. Photolysis expts. were conducted with a Rayonet RPR-200 merry-go-round photoreactor at 253.7 nm. Solns. contg. fixed amt.

of TCE and surfactant Brij 35 were exposed to UV illumination with different concns. of acetone (ACE). Quantum yield in soln. with surfactant Brij 35 and optimum ACE dosage is about 25 times higher than that in Brij 35 alone. However, with an excess ACE dosage, it would act as a light barrier which attenuates the light intensity for TCE photodegrdn. A math. model is therefore developed for the prediction of TCE photodegrdn. in Brij 35 soln. contg. various ACE concns., in which the remaining fraction of TCE (C/C0) in the system can be detd. Apart from the direct photodegrdn., photosensitization is postulated to be another major pathway contributing to the overall decay.

IT 79-01-6, Trichloroethylene, processes  
 (trichloroethene decay in UV system with hydrogen source)  
 RN 79-01-6 HCAPLUS  
 CN Ethene, trichloro- (9CI) (CA INDEX NAME)



IT 9002-92-0, Brij 35  
 (trichloroethene decay in UV system with hydrogen source)  
 RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



CC 61-5 (Water)  
 IT 79-01-6, Trichloroethylene, processes  
 (trichloroethene decay in UV system with hydrogen source)  
 IT 67-64-1, Acetone, reactions 9002-92-0, Brij 35  
 (trichloroethene decay in UV system with hydrogen source)

L71 ANSWER 5 OF 24 HCAPLUS COPYRIGHT 2002 ACS


2001:409205 Document No. 135:199760 The modelling of trichloroethene photodegradation in Brij 35 surfactant by two-stage reaction

. Choy, W. K.; Chu, W. (Department of Civil and Structural Engineering, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong). Chemosphere, 44(2), 211-215 (English) 2001. CODEN: CSMHAF. ISSN: 0045-6535. Publisher: Elsevier Science Ltd..


AB Various clean-up technologies were developed for the removal and/or destruction of trichloroethene (TCE) in the subsurface. Surfactant-aided soil washing followed by photodegrdn. could be a promising approach to such a task. The modeling of TCE photodegrdn. by UV in Brij 35 surfactant micelles is therefore investigated. Two

stages of TCE degrdn. are obsd. in surfactant Brij 35 systems. A lag phase is obsd. at the commencement of the degrdn., but the duration of the lag phase is significantly reduced as the initial pH increases. As the overall decay of TCE is also found to be faster at higher pH levels, it is suggested that the free radical **reaction** is dominant at high pH levels, and the formation of lag phases is mainly due to the deficiency of free radicals at lower pH levels. Since the period of the lag phase gradually decreases with the increase of initial pH level, and the 2 pseudo 1st-order **reaction** consts. (one for the lag phase and one for the subsequent fast decay) for TCE decay in both stages are also pH dependent, a non-steady-state math. model is developed for the prediction of TCE photodegrdn. in Brij 35 solns., in which the remaining fraction of TCE (C/C0) in the system can be detd. at any instant by using a simple parameter of the initial system pH.

- IT 79-01-6, Trichloroethene, **reactions**  
 (modeling of trichloroethene photodegrdn. by UV in Brij 35 surfactant micelles)  
 RN 79-01-6 HCAPLUS  
 CN Ethene, trichloro- (9CI) (CA INDEX NAME)



- IT 9002-92-0, Brij 35  
 (surfactant; modeling of trichloroethene photodegrdn. by UV in Brij 35 surfactant micelles)  
 RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




- CC 60-2 (Waste Treatment and Disposal)  
 Section cross-reference(s): 22, 46  
 IT 79-01-6, Trichloroethene, **reactions**  
 (modeling of trichloroethene photodegrdn. by UV in Brij 35 surfactant micelles)  
 IT 9002-92-0, Brij 35  
 (surfactant; modeling of trichloroethene photodegrdn. by UV in Brij 35 surfactant micelles)


Wing-Ki (Department of Civil and Structural Engineering, Research Centre for Urban Environmental Technology and Management, The Hong Kong Polytechnic University, Hong Kong, Peop. Rep. China). Chemosphere, 41(8), 1199-1204 (English) 2000. CODEN: CSMHAF. ISSN: 0045-6535. Publisher: Elsevier Science Ltd..

AB Photodegrdn. of trichloroethene (TCE) in surfactant micelles was studied in a Rayonet RPR-200 merry-go-round photoreactor at 253.7 nm monochromatic UV lamps, in the presence of surfactants. Surfactants were used as addnl. H sources to improve TCE photodegrdn. rates. About 3 times the rate increment was obsd. in the presence of Brij 35 surfactant micelles vs. water alone. Increasing H<sup>+</sup> and Cl<sup>-</sup> concns. indicated they are the final products of TCE photodegrdn. (i.e., photodechlorination is the dominant mechanism in this system). A lag phase was obsd. at the beginning of the degrdn.; however, the lag phase duration is apparently reduced as initial pH increases. Since overall TCE decay was also faster at higher pH, it is suggested that the free radical **reaction** is dominant at high pH, and formation of lag phases is mainly due to the deficiency of free radicals at lower pH. Photodecompn. of TCE in surfactant micelles also proved to be a clean, effective process; it generates no chlorinated byproducts or intermediates, and TCE is fully decompd. within a reasonable time.

IT 9002-92-0, Brij 35  
(pH effect on lag phase and rate improvement of trichloroethylene photolytic decay in UV/surfactant systems)  
RN 9002-92-0 HCAPLUS  
CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



IT 79-01-6, Trichloroethene, processes  
(pH effect on lag phase and rate improvement of trichloroethylene photolytic decay in UV/surfactant systems)  
RN 79-01-6 HCAPLUS  
CN Ethene, trichloro- (9CI) (CA INDEX NAME)



CC 61-2 (Water)  
Section cross-reference(s): 19, 46, 67  
IT 9002-92-0, Brij 35 9005-65-6, Tween 80  
(pH effect on lag phase and rate improvement of trichloroethylene photolytic decay in UV/surfactant systems)

IT 79-01-6, Trichloroethene, processes  
(pH effect on lag phase and rate improvement of trichloroethylene  
photolytic decay in UV/surfactant systems)

L71 ANSWER 7 OF 24 HCAPLUS COPYRIGHT 2002 ACS

2000:128324 Document No. 132:261508 Classifying environmental  
pollutants: Part 3. External validation of the classification  
system. Verhaar, Henk J. M.; Solbe, John; Speksnijder, John; Van  
Leeuwen, Cees J.; Hermens, Joop L. M. (OpdenKamp, Registration and  
Notification, The Hague, NL-2514 AB, Neth.). Chemosphere, 40(8),  
875-883 (English) 2000. CODEN: CMSHAF. ISSN: 0045-6535.  
Publisher: Elsevier Science Ltd..

AB In order to validate a classification system for the prediction of  
the toxic effect concns. of org. environmental pollutants to fish,  
all available fish acute toxicity data were retrieved from the  
ECETOC database, a database of quality-evaluated aquatic toxicity  
measurements created and maintained by the European Center for the  
Ecotoxicol. and Toxicol. of Chems. The individual chems. for which  
these data were available were classified according to the rule base  
under consideration and predictions of effect concns. or ranges of  
possible effect concns. were generated. These predictions were  
compared to the actual toxicity data retrieved from the database.  
The results of this comparison show that generally the  
classification system provides adequate predictions of either the  
aquatic toxicity (class 1) or the possible range of toxicity (other  
classes) of org. compds. A slight underestimation of effect concns.  
occurs for some highly water-sol., **reactive** chems. with  
low log K<sub>DW</sub> values. On the other end of the scale, some compds.  
that are classified as belonging to a relatively toxic class appear  
to belong to the so-called baseline toxicity compds. For some of  
these, addnl. classification rules are proposed. Furthermore, some  
groups of compds. cannot be classified, although they should be  
amendable to predictions. For these compds. addnl. research as to  
class membership and assocd. prediction rules is proposed.


IT 58-89-9, Lindane 76-44-8, Heptachlor  
79-00-5, 1,1,2-Trichloroethane 79-01-6,  
Trichloroethylene, biological studies 79-34-5,  
1,1,2,2-Tetrachloroethane 120-82-1, 1,2,4-Trichlorobenzene  
3389-71-7 9002-92-0

(external validation of classification system for environmental  
pollutants)


RN 58-89-9. HCAPLUS

CN Cyclohexane, 1,2,3,4,5,6-hexachloro-, (1.alpha.,2.alpha.,3.beta.,4.alpha.,5.alpha.,6.beta.)- (9CI) (CA INDEX NAME)


Relative stereochemistry.




RN 76-44-8 HCAPLUS  
 CN 4,7-Methano-1H-indene, 1,4,5,6,7,8,8-heptachloro-3a,4,7,7a-tetrahydro- (9CI) (CA INDEX NAME)




RN 79-00-5 HCAPLUS  
 CN Ethane, 1,1,2-trichloro- (8CI, 9CI) (CA INDEX NAME)




RN 79-01-6 HCAPLUS  
 CN Ethene, trichloro- (9CI) (CA INDEX NAME)




RN 79-34-5 HCAPLUS  
 CN Ethane, 1,1,2,2-tetrachloro- (8CI, 9CI) (CA INDEX NAME)




RN 120-82-1 HCAPLUS  
 CN Benzene, 1,2,4-trichloro- (8CI, 9CI) (CA INDEX NAME)



RN 3389-71-7 HCAPLUS  
 CN Bicyclo[2.2.1]hepta-2,5-diene, 1,2,3,4,7,7-hexachloro- (9CI) (CA INDEX NAME)



RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



CC 4-3 (Toxicology)  
 Section cross-reference(s): 61  
 IT 51-28-5, 2,4-Dinitrophenol, biological studies 56-38-2, Parathion  
 58-89-9, Lindane 62-53-3, Aniline, biological studies  
 63-25-2, Carbaryl 67-56-1, Methanol, biological studies 67-72-1,



Hexachloroethane 68-12-2, Dimethylformamide, biological studies 71-36-3, 1-Butanol, biological studies 71-43-2, Benzene, biological studies 72-20-8, Endrin 74-90-8, Hydrogen cyanide, biological studies 75-09-2, Dichloromethane, biological studies 75-21-8, Ethylene oxide, biological studies 75-56-9, Propylene oxide, biological studies 76-44-8, Heptachlor 78-92-2, 2-Butanol 79-00-5, 1,1,2-Trichloroethane 79-01-6, Trichloroethylene, biological studies 79-06-1, Acrylamide, biological studies 79-34-5, 1,1,2,2-Tetrachloroethane 80-05-7, Bisphenol A, biological studies 83-32-9, Acenaphthene 84-69-5, 84-74-2, Dibutylphthalate 85-68-7, Benzylbutylphthalate 87-86-5, Pentachlorophenol 88-06-2, 2,4,6-Trichlorophenol 88-30-2, 3-Trifluoromethyl-4-nitrophenol 88-85-7, 2,4-Dinitro-6-(sec-butyl)phenol 95-47-6, o-Xylene, biological studies 95-48-7, o-Cresol, biological studies 95-57-8, 2-Chlorophenol 96-22-0, 3-Pentanone 98-51-1, 4-tert-Butyltoluene 100-02-7, 4-Nitrophenol, biological studies 100-42-5, Styrene, biological studies 100-52-7, Benzaldehyde, biological studies 105-67-9, 2,4-Dimethylphenol 106-42-3, p-Xylene, biological studies 106-44-5, p-Cresol, biological studies 106-46-7, 1,4-Dichlorobenzene 106-51-4, 2,5-Cyclohexadiene-1,4-dione, biological studies 106-68-3, 3-Octanone 106-89-8, Epichlorohydrin, biological studies 106-92-3, Allylglycidyl ether 107-02-8, Acrolein, biological studies 107-05-1, Allyl chloride 107-07-3, 2-Chloroethanol, biological studies 107-18-6, Allyl alcohol, biological studies 107-70-0, 108-10-1, Methyl isobutyl ketone 108-20-3, Diisopropyl ether 108-38-3, m-Xylene, biological studies 108-39-4, m-Cresol, biological studies 108-68-9, 3,5-Dimethylphenol 108-88-3, Toluene, biological studies 108-90-7, Chlorobenzene, biological studies 108-95-2, Phenol, biological studies 109-70-6, 1-Chloro-3-bromopropane 110-49-6, Ethyleneglycol monomethyl ether acetate 111-15-9, Ethyleneglycol monoethyl ether acetate 111-78-4, 1,5-Cyclooctadiene 111-87-5, 1-Octanol, biological studies 112-30-1, Decanol 112-34-5, Diethyleneglycol monobutyl ether 115-29-7, Endosulfan 115-32-2, Kelthane 116-06-3, Aldicarb 118-79-6, 2,4,6-Tribromophenol 120-82-1, 1,2,4-Trichlorobenzene 120-83-2, 2,4-Dichlorophenol 121-46-0, Bicyclo(2.2.1)hepta-2,5-diene 121-75-5, Malathion 122-60-1, Phenylglycidyl ether 123-54-6, 2,4-Pentanedione, biological studies 127-18-4, Tetrachloroethylene, biological studies 141-79-7, Mesityl oxide 142-28-9, 1,3-Dichloropropane 143-08-8, 1-Nonanol 298-00-0, Methylparathion 298-04-4, Disulfoton 333-41-5, Diazinon 504-20-1, Phorone 534-52-1, 2-Methyl-4,6-dinitrophenol 544-25-2, 1,3,5-Cycloheptatriene 563-47-3, 3-Chloro-2-methylpropene 576-26-1, 2,6-Dimethylphenol 577-19-5, 2-Bromonitrobenzene 627-30-5, 3-Chloropropanol 693-21-0, Diethyleneglycol dinitrate 935-95-5, 2,3,5,6-Tetrachlorophenol 944-22-9, Fonofos 1204-21-3, 1471-17-6, Pentaerythritol triallyl ether 1582-09-8, Trifluralin 1912-24-9, Atrazine 1918-02-1, Picloram 1962-75-0, Dibutylterephthalate 2212-67-1, Molinate 2461-15-6, 2-Ethylhexylglycidyl ether 2921-88-2, Chlorpyrifos

3389-71-7 3698-83-7, 4,6-Dichloro-1,3-dinitrobenzene  
 4904-61-4, 1,5,9-Cyclododecatriene 6515-38-4, 3,5,6-Trichloro-2-  
 pyridinol 9002-92-0 9002-93-1, Triton X-45 9016-45-9  
 11097-69-1, PCB 1254 28249-77-6, Thiobencarb 31557-34-3,  
 2-Methoxy-3,5,6-trichloropyridine 51630-58-1, Fenvalerate  
 52645-53-1, Permethrin 52918-63-5, Deltamethrin 53469-21-9, PCB  
 1242 66330-88-9, Hydrothol-191 70124-77-5, AC 222705  
 (external validation of classification system for environmental  
 pollutants)

L71 ANSWER 8 OF 24 HCAPLUS COPYRIGHT 2002 ACS


1999:589104 Document No. 131:310372 Study of the **Reaction**  
 1,1,1-Trichloro-2,2-bis(p-chlorophenyl)ethane (DDT) + OH<sup>-</sup> in Nonionic  
 Micellar Solutions. Munoz, Maria; Rodriguez, Amalia; Graciani,  
 Maria del Mar; Ortega, Francisco; Vazquez, Maria; Moya, Maria Luisa  
 (Departamento de Quimica Fisica, Universidad de Sevilla, Seville,  
 41012, Spain). Langmuir, 15(22), 7876-7879 (English) 1999. CODEN:  
 LANGD5. ISSN: 0743-7463. Publisher: American Chemical Society.

AB Kinetic micellar effects for the title **reaction** in aq.  
 Brij35 and Triton X-100 nonoionic micellar solns. were rationalized  
 by structural studies using surface tension, as well as fluorescence  
 and light scattering measurements.

IT 9002-92-0, Brij35  
 (kinetic micellar effects in dehydrochlorination of DDT in  
 presence of hydroxide in aq.)

RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




IT 50-29-3, DDT, **reactions**

(kinetic micellar effects in dehydrochlorination of DDT in  
 presence of hydroxide in aq.)

RN 50-29-3 HCAPLUS

CN Benzene, 1,1'-(2,2,2-trichloroethylidene)bis[4-chloro- (9CI) (CA  
 INDEX NAME)



CC 22-13 (Physical Organic Chemistry)

Section cross-reference(s): 66

- IT 9002-92-0, Brij35 9002-93-1, Triton X-100  
(kinetic micellar effects in dehydrochlorination of DDT in presence of hydroxide in aq.)
- IT 50-29-3, DDT, **reactions**  
(kinetic micellar effects in dehydrochlorination of DDT in presence of hydroxide in aq.)
- IT 14280-30-9, Hydroxide, **reactions**  
(kinetic micellar effects in dehydrochlorination of DDT in presence of hydroxide in aq.)

L71 ANSWER 9 OF 24 HCAPLUS COPYRIGHT 2002 ACS

1998:793401 Document No. 130:85409 Photodechlorination Mechanism of DDT in a UV/Surfactant System. Chu, Wei (Department of Civil and Structural Engineering, Hong Kong Polytechnic University, Kowloon, Hong Kong). Environmental Science and Technology, 33(3), 421-425 (English) 1999. CODEN: ESTHAG. ISSN: 0013-936X. Publisher: American Chemical Society.


AB The photochem. **reactions** of the organochlorine pesticide DDT in aq. solns. contg. nonionic surfactant micelles (Brij 35, Brij 52, and Brij 72) were studied and modeled. All photolytic expts. were conducted in a Rayonet RPR-200 merry-go-round photoreactor using a 253.7-nm Hg monochromatic UV lamps. Pseudo-1st-order decay through photodechlorination was the dominant **reaction** pathway for DDT photodecay. The primary photoproducts include lesser chlorinated compds. (DDE and DDD) and HCl. The photodechlorination of DDT involves 2 stages; the 1st is the fast aliph. Cl redn., followed by a slow arom. Cl redn. The photodecay rates of DDT were doubled in the Brij 52 micellar soln. compared to that in water alone. A 1st-order parallel/consecutive model was developed to predict the photodecay of DDT and the generation of DDE/DDD in the micellar/aq. system.

IT 72-54-8, Ddd 72-55-9, DDE, formation  
(nonpreparative)

(photodechlorination mechanism of DDT in UV/surfactant system)


RN 72-54-8 HCAPLUS

CN Benzene, 1,1'-(2,2-dichloroethylenidene)bis[4-chloro- (9CI) (CA INDEX NAME)




RN 72-55-9 HCAPLUS


CN Benzene, 1,1'-(dichloroethenylenidene)bis[4-chloro- (9CI) (CA INDEX NAME)




IT 9002-92-0, Brij 35 9004-95-9, Brij 52  
 9005-00-9, Brij 72  
 (photodechlorination mechanism of DDT in UV/surfactant system)  
 RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




RN 9004-95-9 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



RN 9005-00-9 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-octadecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)



IT 50-29-3, DDT, processes  
 (photodechlorination mechanism of DDT in UV/surfactant system)  
 RN 50-29-3 HCAPLUS  
 CN Benzene, 1,1'-(2,2,2-trichloroethylidene)bis[4-chloro- (9CI) (CA  
 INDEX NAME)




- CC 60-2 (Waste Treatment and Disposal)  
Section cross-reference(s): 5
- IT 72-54-8, Ddd 72-55-9, DDE, formation  
(nonpreparative) 7647-01-0, Hydrogen chloride, formation  
(nonpreparative)  
(photodechlorination mechanism of DDT in UV/surfactant system)
- IT 9002-92-0, Brij 35 9004-95-9, Brij 52  
9005-00-9, Brij 72  
(photodechlorination mechanism of DDT in UV/surfactant system)
- IT 50-29-3, DDT, processes  
(photodechlorination mechanism of DDT in UV/surfactant system)
- L71 ANSWER 10 OF 24 HCAPLUS COPYRIGHT 2002 ACS  
1998:325409 Document No. 129:19220 Phototransformations of  
Polychlorobiphenyls in Brij 58 Micellar Solutions. Chu, Wei;  
Jafvert, Chad T.; Diehl, Claude A.; Marley, Karen; Larson, Richard  
A. (School of Civil Engineering, Purdue University, West Lafayette,  
IN, 47907-1284, USA). Environmental Science and Technology, 32(13),  
1989-1993 (English) 1998. CODEN: ESTHAG. ISSN: 0013-936X.  
Publisher: American Chemical Society.
- AB Our purpose in conducting these studies was to examine photolysis as  
a destructive process for polychlorobiphenyls (PCBs) extd. from  
soils with surfactant solns. Surfactants have shown promise as  
agents for removing free-phase and sorbed contaminants from soils,  
yet information on ultimate disposal options and recycle/recovery  
strategies for the surfactants is generally lacking. For aryl  
halides, photodechlorination may result in decontamination,  
eliminating the need to phys. sep. these contaminants from the  
washing soln. Photochem. **reactions** of the PCB congener  
mixt., Aroclor 1254, and the specific congener, 2,3,4,5-  
tetrachlorobiphenyl (2,3,4,5-TeCB), were investigated in aq. solns.  
contg. surfactant micelles with UV light at 253.7 nm. Photoredn.  
through photodechlorination was shown to be the main decay pathway  
in which lesser chlorinated congeners were formed as intermediates.  
In expts. with 2,3,4,5-TeCB, final noncarbon-contg. products  
included Cl<sup>-</sup> and H<sup>+</sup>, both produced nearly stoichiometrically from  
the starting materials. The quantum yield for decay of 0.1.μM  
2,3,4,5-TeCB in 0.5 mM Brij 58 micellar solns. was over six times  
greater than in water alone. Sequential extn. from a soil and  
photoredn. of 2,3,4,5-tetrachlorobiphenyl by Brij 58 solns. proved  
to be limited by surfactant loss to the soil.
- IT 9004-95-9, Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-  
.omega.-hydroxy-

(phototransformations of polychlorobiphenyls in Brij 58 micellar solns. in relation to soil remediation)

RN 9004-95-9 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)




IT 33284-53-6, 2,3,4,5-Tetrachlorobiphenyl

(phototransformations of polychlorobiphenyls in Brij 58 micellar solns. in relation to soil remediation)

RN 33284-53-6 HCAPLUS

CN 1,1'-Biphenyl, 2,3,4,5-tetrachloro- (9CI) (CA INDEX NAME)



CC 60-4 (Waste Treatment and Disposal)

Section cross-reference(s): 19, 46, 74

IT 9004-95-9, Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy-

(phototransformations of polychlorobiphenyls in Brij 58 micellar solns. in relation to soil remediation)

IT 92-52-4D, 1,1'-Biphenyl, chloro derivs., **reactions**

11097-69-1, Aroclor 1254 33284-53-6, 2,3,4,5-Tetrachlorobiphenyl

(phototransformations of polychlorobiphenyls in Brij 58 micellar solns. in relation to soil remediation)

L71 ANSWER 11 OF 24 HCAPLUS COPYRIGHT 2002 ACS

1998:137232 Document No. 128:267091 Induction of forward mutations at the thymidine kinase locus of mouse lymphoma cells: evidence for electrophilic and non-electrophilic mechanisms. Henry, B.; Grant, S. G.; Klopman, G.; Rosenkranz, H. S. (260 Kappa Drive, Department of Environmental and Occupational Health, University of Pittsburgh, Pittsburgh, PA, 15238, USA). Mutation Research, 397(2), 313-335 (English) 1998. CODEN: MUREAV. ISSN: 0027-5107. Publisher: Elsevier Science B.V..

AB A database of 209 chems. tested for induction of forward mutations at the heterozygous thymidine kinase (TK.+-. ) locus in L5178Y mouse lymphoma cells was analyzed for structure-activity relationships


using the MultiCASE expert system. Consistent with evidence of several contributing biol. mechanisms, these studies suggest that such mutations may occur by more than one mechanism. As might be expected, there was evidence for a component involving direct electrophilic attack on the cellular DNA, in a manner previously established as causative in the induction of mutations in Salmonella. In addn., however, there was also strong evidence for another mechanism or mechanisms involving chromosome missegregation, cellular toxicity or an alternate site of action, such as the microtubules.

IT 72-55-9, biological studies 75-25-2, Bromoform  
76-01-7, Pentachloroethane 76-44-8, Heptachlor  
79-01-6, Trichloroethylene, biological studies  
630-20-6 3322-93-8, 1-(1,2-Dibromoethyl)-3,4-  
dibromocyclohexane 9002-92-0

(forward mutations induction at the thymidine kinase locus of mouse lymphoma cells - evidence for electrophilic and non-electrophilic mechanisms)


RN 72-55-9 HCAPLUS

CN Benzene, 1,1'-(dichloroethenylidene)bis[4-chloro- (9CI) (CA INDEX NAME)




RN 75-25-2 HCAPLUS

CN Methane, tribromo- (8CI, 9CI) (CA INDEX NAME)




RN 76-01-7 HCAPLUS

CN Ethane, pentachloro- (8CI, 9CI) (CA INDEX NAME)




RN 76-44-8 HCAPLUS

CN 4,7-Methano-1H-indene, 1,4,5,6,7,8,8-heptachloro-3a,4,7,7a-tetrahydro- (9CI) (CA INDEX NAME)




RN 79-01-6 HCAPLUS

CN Ethene, trichloro- (9CI) (CA INDEX NAME)




RN 630-20-6 HCAPLUS

CN Ethane, 1,1,1,2-tetrachloro- (8CI, 9CI) (CA INDEX NAME)



RN 3322-93-8 HCAPLUS

CN Cyclohexane, 1,2-dibromo-4-(1,2-dibromoethyl)- (7CI, 8CI, 9CI) (CA INDEX NAME)




RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)



(CA INDEX NAME)



CC 4-6 (Toxicology)

Section cross-reference(s): 3, 7

IT DNA

(**reactivity**; forward mutations induction at the thymidine kinase locus of mouse lymphoma cells - evidence for electrophilic and non-electrophilic mechanisms)


IT 50-55-5, Reserpine 50-81-7, L-Ascorbic acid, biological studies  
 51-03-6, Piperonyl butoxide 51-79-6, Urethane 52-68-6,  
 Trichlorfon 53-96-3, 2-Acetylaminofluorene 54-31-9, Furosemide  
 54-85-3, Isoniazid 55-86-7, Nitrogen mustard 56-49-5,  
 3-Methylcholanthrene 56-57-5, 4-Nitroquinoline-1-oxide 56-72-4,  
 Coumaphos 57-50-1, Sucrose, biological studies 57-97-6,  
 9,10-Dimethyl-1,2-benzanthracene 58-93-5, Hydrochlorothiazide  
 59-42-7, Phenylephrine 59-87-0, Nitrofurazone 60-54-8,  
 Tetracycline 60-57-1, Dieldrin 61-82-5, 3-Aminotriazole  
 62-50-0, Ethyl methanesulfonate 62-53-3, Aniline, biological  
 studies 62-73-7, Dichlorvos 65-61-2, Acridine orange 66-27-3,  
 Methyl methanesulfonate 67-20-9, Nitrofurantoin 68-12-2,  
 Dimethylformamide, biological studies 69-65-8, Mannitol 69-74-9,  
 Cytarabine hydrochloride 70-25-7, N-Methyl-N'-nitro-N-  
 nitrosoguanidine 71-43-2, Benzene, biological studies 72-20-8,  
 Endrin 72-55-9, biological studies 75-09-2, Methylene  
 chloride, biological studies 75-25-2, Bromoform 75-27-4,  
 Bromodichloromethane 75-35-4, Vinylidene chloride, biological  
 studies 75-56-9, Propylene oxide, biological studies 75-65-0,  
 tert-Butyl alcohol, biological studies 76-01-7,  
 Pentachloroethane 76-44-8, Heptachlor 78-42-2,  
 Tris(2-ethylhexyl)phosphate 78-59-1, Isophorone 78-87-5  
 79-01-6, Trichloroethylene, biological studies 79-11-8,  
 Chloracetic acid, biological studies 79-57-2, Oxytetracycline  
 80-05-7, Bisphenol A, biological studies 80-08-0,  
 4,4'-Sulfonyldianiline 80-62-6, Methyl methacrylate 83-79-4,  
 Rotenone 85-68-7, Butyl benzyl phthalate 86-30-6,  
 Diphenylnitrosamine 86-73-7, Fluorene 87-29-6, Cinnamyl  
 anthranilate 88-06-2, 2,4,6-Trichlorophenol 88-96-0, Phthalamide  
 89-25-8, 1-Phenyl-3-methyl-5-pyrazolone 89-78-1 91-08-7,  
 2,6-Toluene diisocyanate 91-22-5, Quinoline, biological studies  
 91-80-5, Methapyrilene 91-81-6, Tripelennamine 92-52-4D,  
 Biphenyl, bromo derivs. 92-87-5, Benzidine 92-93-3,  
 4-Nitrobiphenyl 94-20-2, Chlorpropamide 94-59-7, Safrole  
 95-50-1, 1,2-Dichlorobenzene 95-51-2, o-Chloroaniline 95-53-4,  
 o-Toluidine, biological studies 95-79-4, 5-Chloro-o-toluidine  
 95-86-3, 2,4-Diaminophenol 96-12-8 96-31-1, N,N'-Dimethylurea  
 96-45-7, N,N'-Ethylenethiourea 97-53-0, Eugenol 97-77-8,

Disulfiram 98-01-1, Furfural, biological studies 99-56-9,  
4-Nitro-o-phenylenediamine 99-57-0, 2-Amino-4-nitrophenol  
100-02-7, p-Nitrophenol, biological studies 100-41-4, Ethyl  
benzene, biological studies 100-44-7, Benzyl chloride, biological  
studies 100-51-6, Benzyl alcohol, biological studies 100-52-7,  
Benzaldehyde, biological studies 101-77-9, 4,4'-Methylenedianiline  
101-80-4, 4,4'-Oxydianiline 101-90-6, Diglycidyl resorcinol ether  
103-23-1, Di(2-ethylhexyl)adipate 105-11-3, p-Benzoquinone dioxime  
105-55-5, 1,3-Diethyl-2-thiourea 105-87-3, Geranyl acetate  
106-40-1, p-Bromoaniline 106-46-7, 1,4-Dichlorobenzene 106-47-8,  
p-Chloroaniline, biological studies 106-88-7 106-99-0,  
1,3-Butadiene, biological studies 107-07-3, Ethylene chlorohydrin,  
biological studies 107-21-1, Ethylene glycol, biological studies  
108-46-3, Resorcinol, biological studies 108-60-1,  
Bis(2-chloro-1-methylethyl)ether 108-78-1, Melamine, biological  
studies 108-88-3, Toluene, biological studies 108-90-7,  
Chlorobenzene, biological studies 108-94-1, Cyclohexanone,  
biological studies 108-95-2, Phenol, biological studies  
109-69-3, N-Butyl chloride 110-00-9, Furan 110-86-1, Pyridine,  
biological studies 111-30-8, Glutaraldehyde 115-07-1, Propylene,  
biological studies 115-28-6, Chlorendic acid 115-29-7,  
Endosulfan 120-61-6, Dimethyl terephthalate 120-83-2,  
2,4-Dichlorophenol 121-79-9, Propyl gallate 121-88-0,  
2-Amino-5-nitrophenol 123-30-8 123-31-9, Hydroquinone,  
biological studies 123-91-1, 1,4-Dioxane, biological studies  
124-48-1, Chlorodibromomethane 124-64-1 126-92-1 127-18-4,  
Tetrachloroethylene, biological studies 127-69-5, Sulfisoxazole  
128-37-0, BHT, biological studies 131-17-9, Diallyl phthalate  
133-06-2, Captan 135-88-6 136-40-3, Phenazopyridine  
hydrochloride 136-77-6, 4-Hexylresorcinol 137-30-4, Ziram  
140-11-4, Benzyl acetate 140-88-5, Ethyl acrylate 142-28-9,  
1,3-Dichloropropane 148-18-5, Sodium diethyldithiocarbamate  
148-24-3, 8-Hydroxyquinoline, biological studies 149-30-4,  
2-Mercaptobenzothiazole 150-38-9, Trisodium EDTA 150-68-5,  
Monuron 151-21-3, SDS, biological studies 271-89-6,  
2,3-Benzofuran 298-18-0 320-67-2, 5-Azacytidine 333-41-5,  
Diazinon 366-70-1, Natulan 434-13-9, Lithocholic acid  
505-60-2, Mustard gas 510-15-6, Chlorobenzilate 513-37-1,  
.beta.,.beta.-Dimethylvinyl chloride 518-47-8, Acid yellow 73  
542-75-6, 1,3-Dichloropropene 542-78-9, Malonaldehyde 563-47-3,  
3-Chloro-2-methyl-1-propene 569-61-9, C.I. Basic red 9 584-84-9  
597-25-1, Dimethyl morpholinophosphoramidate 598-55-0, Methyl  
carbamate 607-57-8, 2-Nitrofluorene 609-20-1,  
2,6-Dichloro-p-phenylenediamine 624-83-9, Methyl isocyanate  
630-20-6 706-87-6 842-07-9, C.I. Solvent yellow 14  
1163-19-5 1212-29-9, 1,3-Bis(cyclohexyl)thiourea 1239-45-8,  
Ethidium bromide 1596-84-5, Succinic acid 2,2-dimethylhydrazide  
1634-78-2, Malaoxone 1746-01-6, TCDD 1825-21-4,  
Pentachloroanisole 1897-45-6 1910-42-5, Methyl viologen  
1936-15-8 2164-17-2, Fluometuron 2185-92-4, 2-Biphenylamine  
hydrochloride 2432-99-7 2438-88-2, 2,3,5,6-Tetrachloro-4-  
nitroanisole 2475-45-8, C.I. Disperse blue 1 2489-77-2


2783-94-0, FD and C yellow 6 2784-94-3, HC blue no. 1 2832-40-8,  
 C.I. Disperse yellow 3 3105-97-3, Hycanthone 3131-60-0,  
 6-Azacytidine 3322-93-8, 1-(1,2-Dibromoethyl)-3,4-  
 dibromocyclohexane 3546-10-9, Phenesterine 3567-69-9, C.I. Acid  
 red 14 4460-86-0, 2,4,5-Trimethoxy benzaldehyde 5160-02-1  
 5307-14-2, 2-Nitro-p-phenylenediamine 5989-27-5, D-Limonene  
 6959-48-4, 3-Chloromethylpyridine hydrochloride 7177-48-2,  
 Ampicillin trihydrate 7320-37-8, 1,2-Epoxyhexadecane  
 9002-92-0 12789-03-6, Chlordane 15481-70-6,  
 2,6-Toluenediamine dihydrochloride 17924-92-4, Zearalenone  
 18883-66-4, Streptozotocin 21739-91-3, Cytembena 28322-02-3,  
 N-4-Fluorenylacetamide 33229-34-4, HC blue no. 2 41372-08-1  
 54150-69-5, 2,4-Dimethoxyaniline hydrochloride 65589-70-0,  
 Acriflavine

(forward mutations induction at the thymidine kinase locus of  
 mouse lymphoma cells - evidence for electrophilic and  
 non-electrophilic mechanisms)


- L71 ANSWER 12 OF 24 HCAPLUS COPYRIGHT 2002 ACS  
 1996:379316 Document No. 125:34453 Recovery of epoxidized block  
 copolymers by steam stripping. Oshino, Yasuhiro; Ootsuka, Yoshihiro  
 (Daicel Chem, Japan). Jpn. Kokai Tokkyo Koho JP 08059734 A2  
 19960305 Heisei, 7 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP  
 1994-191612 19940815.
- AB Slurries or solns. of epoxidized (hydrogenated) arom. vinyl  
 compd.-conjugated diene block copolymers in org. solvents are  
 subjected to steam stripping in the presence of surfactants to  
 recover the polymers. Thus, a soln. of epoxidized TR 2000  
 (styrene-butadiene-styrene block copolymer) (I) in AcOEt was  
 subjected to steam stripping in the presence of Emulgen PP 290  
 (nonionic surfactant) at 90.degree. to recover I without adhesion to  
 the inner wall of the reactor or the stirring paddle.
- IT 67-66-3, Chloroform, uses 9002-92-0, Emulgen 147  
 (surfactants in recovery of epoxidized block copolymers in org.  
 solvents by steam stripping)
- RN 67-66-3 HCAPLUS  
 CN Methane, trichloro- (9CI) (CA INDEX NAME)



- RN 9002-92-0 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




- IC ICM C08F006-00  
ICS C08F236-10; C08G059-34
- CC 35-10 (Chemistry of Synthetic High Polymers)
- IT 67-66-3, Chloroform, uses 108-88-3, Toluene, uses 110-54-3, Hexane, uses 110-82-7, Cyclohexane, uses 112-00-5, Quartamin 24P 141-78-6, Ethyl acetate, uses 1330-20-7, Xylene, uses 9002-92-0, Emulgen 147 9003-11-6, Emulgen PP 290 9004-98-2, Emulgen 408 9016-45-9, Emulgen 985 88984-51-4, Electrostripper F  
(surfactants in recovery of epoxidized block copolymers in org. solvents by steam stripping)
- L71 ANSWER 13 OF 24 HCAPLUS COPYRIGHT 2002 ACS  
1995:343404 Document No. 122:169044 Biochemical degradability of selected organic compounds; part 2. Wotzka, von Joerg; Pfitzner, Steffi; Giest, Baerbel (Bundesanstalt fur Gewasserkunde, Aussenstelle Berlin, Berlin, 12439, Germany). Deutsche Gewaesserkundliche Mitteilungen, 38(1-2), 10-17 (German) 1994. CODEN: DGMTAO. ISSN: 0012-0235.
- AB The biodegradability of a substance is a relevant characteristic to assess its behavior in wastewater treatment, its accumulation in the environment, and the water pollution hazard it poses. Dissimilation of org. compds. was examd. in lab.-scale wastewater treatment plants and in respirometers (Sapromat, Warburg app.) by adapted and non-adapted microorganisms. In evaluating the results, substances were assigned to 1 of 4 groups (A-D) on the basis of the BOD:COD ratio or calcd. theor. BOD. A total of 178 selected substances were tested; results are summarized. These 4 groups were established on the basis of the chem. structure of the compds. A total of 20 figures display degrdn. curves measured by respirometer, reflecting the effect of adaptation, dry sludge content, **reaction** time, and delayed dissimilation. Distribution of the tested chems. among the 4 groups is: biochem. easily degradable (27 substances, 15%); biochem. degradable (60 substances, 34%); biochem. hardly degradable (11 substances, 6%); and non-biodegradable (80 substances, 45%). Alcs., esters, org. acids, phenooxyacids, and surfactants belong to the biochem. easily degradable or biochem. degradable substances, Groups A and B, whereas org. Sn compds., sulfonic acids, triazine, and most pesticides were among the non-biodegradable compds.
- IT 98-07-7, Benzotrichloride 27306-79-2  
(biochem. degrdn. org. compds. in lab.-scale wastewater treatment plants and respirometers by adapted and non-adapted microorganisms)
- RN 98-07-7 HCAPLUS
- CN Benzene, (trichloromethyl)- (9CI) (CA INDEX NAME)



RN 27306-79-2 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-tetradecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



CC 60-2 (Waste Treatment and Disposal)

Section cross-reference(s): 5, 45, 46, 61

IT 50-33-9, Phenylbutazone, biological studies 51-28-5,  
2,4-Dinitrophenol, biological studies 52-51-7, Bronopol 59-50-7,  
p-Chloro-m-cresol 60-51-5, Dimethoate 61-82-5, Amitrol  
62-73-7, Dichlorvos 63-25-2, Carbaryl 70-30-4, Hexachlorophene  
75-87-6, Chloral 81-11-8, Flavonic acid 93-65-2,  
2-(4-Chloro-2-methylphenoxy)-propionic acid 94-74-6,  
4-Chloro-2-methyl-phenoxyacetic acid 94-75-7, 2,4-  
Dichlorophenoxyacetic acid, biological studies 94-81-5,  
4-(4-Chloro-2-methylphenoxy)butyric acid 94-82-6,  
4-(2,4-Dichloro-phenoxy)butyric acid 95-55-6, o-Aminophenol  
95-57-8, o-Chlorophenol 96-91-3, 4,6-Dinitro-2-aminophenol  
**98-07-7**, Benzotrichloride 98-11-3D, Benzene sulfonic acid,  
alkyl derivs. 98-47-5, m-Nitrobenzene sulfonic acid 98-95-3,  
biological studies 99-57-0, 2-Amino-4-nitrophenol 100-02-7,  
p-Nitrophenol, biological studies 100-17-4 101-21-3,  
Chlorpropham 101-42-8, Fenuron 103-90-2, 4-Acetaminophenol  
104-74-5, Dodecylpyridinium chloride 106-48-9, p-Chlorophenol  
108-77-0, Cyanurichloride 108-90-7, Chlorobenzene, biological  
studies 108-94-1, Cyclohexanone, biological studies 110-82-7,  
Cyclohexane, biological studies 111-30-8, Glutardialdehyde  
116-52-9, Dichloral urea 119-19-7, Phenyl .gamma.-acid 120-36-5,  
2-(2,4-Dichlorophenoxy)-propionic acid 120-83-2,  
2,4-Dichlorophenol 121-47-1, m-Aminobenzene sulfonic acid  
121-57-3, p-Aminobenzene sulfonic acid 122-34-9, Simazine  
122-42-9, Proptham 122-87-2, p-Hydroxyphenylglycine 123-30-8,  
p-Aminophenol 127-06-0 127-81-1, Solupront 135-19-3,  
2-Naphthalenol, biological studies 137-26-8, Thiuram 139-40-2,  
Propazine 149-30-4, 2(3H)-Benzothiazolethione 150-68-5, Monuron  
287-92-3, Cyclopentane 298-00-0, Parathionmethyl 330-54-1,  
Diuron 330-55-2, Linuron 480-96-6, Benzofuroxan 498-66-8,  
Norbornene 534-52-1 591-27-5, m-Aminophenol 645-62-5

705-62-4 834-12-8, Ametryn 873-50-7, 1-Carbamoyl-3-methylpyrazole 973-21-7, Dinobuton 999-81-5 1007-36-9, Defenuron 1014-69-3, Desmetryn 1570-64-5, p-Chloro-o-cresol 1689-83-4, Ioxynil 1689-84-5, Bromoxynil 1712-64-7, Isopropylnitrate 1746-81-2, Monolinuron 1836-75-5, Nitrofen 1912-24-9, Atrazine 2164-08-1, Lenacil 2828-42-4, Proximpham 3060-89-7, Metobromuron 3408-97-7, Bromfenuron 4499-99-4, Triethylene glycol diethylether 5234-68-4, Carboxin 5629-51-6, Dinobutonmethyl 6920-97-4, 1,2-Dimethyl-5-vinylpyridinium methylsulfate 7287-19-6, Prometryn 7398-69-8, Dimethyl-diallyl-ammonium chloride 7540-60-5, Cyclohexylsulfamate 9016-45-9 10265-92-6, Methamidophos 13684-63-4, Phenmedipham 19961-72-9 25154-42-1, Chlorobutane 25366-23-8, Thiazafuron 26377-90-2 **27306-79-2** 28730-17-8, Methfuroxam 29756-37-4, Chloroheptane 39549-27-4 66719-08-2 78846-76-1 88265-20-7, Trizin 91315-15-0, Aldimorph 161544-84-9, Tridiron (biochem. degrdn. org. compds. in lab.-scale wastewater treatment plants and respirometers by adapted and non-adapted microorganisms)

L71 ANSWER 14 OF 24 HCAPLUS COPYRIGHT 2002 ACS


1994:662596 Document No. 121:262596 Photodechlorination of Polychlorobenzene Congeners in Surfactant Micelle Solutions. Chu, Wei; Jafvert, Chad T. (School of Civil Engineering, Purdue University, West Lafayette, IN, 47907, USA). Environmental Science and Technology, 28(13), 2415-22 (English) 1994. CODEN: ESTHAG. ISSN: 0013-936X.

AB Photochem. **reactions** of polychlorobenzene congeners in aq. solns. contg. surfactant micelles were investigated. All photolysis expts. were performed with a Rayonet RPR-100 merry-go-round photoreactor utilizing 253.7-nm monochromatic UV lamps. Photoredn. through photodechlorination was the main decay pathway in which lesser chlorinated congeners and benzene were formed as intermediates. Final products included H<sup>+</sup> and Cl<sup>-</sup> in approx. stoichiometric amts. In addn., some minor pathways were obsd. including photochlorination (the reverse **reaction**), photoisomerization, and finally photohydrolysis, through which phenol was found. The quantum yield for the decay of hexachlorobenzene was about an order of magnitude greater in micellar solns. than in water alone. Several H sources were investigated with NaBH<sub>4</sub> and is a promising rate enhancer at elevated concns.

IT 9002-92-0, Brij 35 9004-95-9, Brij 58  
(photodechlorination of polychlorobenzene congeners in surfactant micelle solns.)


RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



RN 9004-95-9 HCAPLUS


CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



IT 87-61-6, 1,2,3-Trichlorobenzene 95-94-3,  
1,2,4,5-Tetrachlorobenzene 108-70-3, 1,3,5-  
Trichlorobenzene 120-82-1, 1,2,4-Trichlorobenzene  
608-93-5, Pentachlorobenzene 634-66-2,  
1,2,3,4-Tetrachlorobenzene 634-90-2, 1,2,3,5-  
Tetrachlorobenzene  
(photodechlorination of polychlorobenzene congeners in surfactant  
micelle solns.)


RN 87-61-6 HCAPLUS

CN Benzene, 1,2,3-trichloro- (8CI, 9CI) (CA INDEX NAME)




RN 95-94-3 HCAPLUS

CN Benzene, 1,2,4,5-tetrachloro- (8CI, 9CI) (CA INDEX NAME)




RN 108-70-3 HCAPLUS


CN Benzene, 1,3,5-trichloro- (8CI, 9CI) (CA INDEX NAME)




RN 120-82-1 HCAPLUS  
CN Benzene, 1,2,4-trichloro- (8CI, 9CI) (CA INDEX NAME)



RN 608-93-5 HCAPLUS  
CN Benzene, pentachloro- (6CI, 8CI, 9CI) (CA INDEX NAME)




RN 634-66-2 HCAPLUS  
CN Benzene, 1,2,3,4-tetrachloro- (7CI, 8CI, 9CI) (CA INDEX NAME)



RN 634-90-2 HCAPLUS  
CN Benzene, 1,2,3,5-tetrachloro- (7CI, 8CI, 9CI) (CA INDEX NAME)






- CC 60-2 (Waste Treatment and Disposal)  
Section cross-reference(s): 61
- IT 9002-92-0, Brij 35 9004-95-9, Brij 58 9005-64-5,  
Tween 20 9005-65-6, Tween 80  
(photodechlorination of polychlorobenzene congeners in surfactant  
micelle solns.)
- IT 87-61-6, 1,2,3-Trichlorobenzene 95-50-1,  
1,2-Dichlorobenzene 95-94-3, 1,2,4,5-Tetrachlorobenzene  
106-46-7, 1,4-Dichlorobenzene 108-70-3,  
1,3,5-Trichlorobenzene 108-90-7, Monochlorobenzene, occurrence  
118-74-1, Hexachlorobenzene 120-82-1, 1,2,4-  
Trichlorobenzene 541-73-1, 1,3-Dichlorobenzene 608-93-5,  
Pentachlorobenzene 634-66-2, 1,2,3,4-Tetrachlorobenzene  
634-90-2, 1,2,3,5-Tetrachlorobenzene  
(photodechlorination of polychlorobenzene congeners in surfactant  
micelle solns.)
- L71 ANSWER 15 OF 24 HCAPLUS COPYRIGHT 2002 ACS  
1993:549321 Document No. 119:149321 Effect of substituents and  
environment on the photochromic properties of spiropyrans. Tomioka,  
Hideo; Zhao, Xiutai (Fac. Eng., Mie Univ., Tsu, 514, Japan). Nippon  
Kagaku Kaishi (7), 884-90 (Japanese) 1993. CODEN: NKAKB8. ISSN:  
0369-4577.
- AB 1',3',3'-Trimethyl-6-nitrospiro[2H-1-benzopyran-2,2'-indolines]  
(SPs) having methoxy groups on the indoline ring were prepd. and  
their photochromic properties were investigated in org. solvents and  
in films. SPs showed normal photochromism in the less polar  
solvents, while reverse photochromism occurred as the solvent  
polarity increased. Linear relationship between electronic  
transition energy (EMC) of the colored species and log k was obsd.  
(k = the rate const. for coloration or decoloration process). In  
micellar soln.,  $\lambda_{max}$  underwent significant blue shift and no  
linear relationship between EMC and log k was obsd. In the films of  
PMMA or polyion complexes of poly(styrene sulfonate) and  
distearyldimethylammonium chloride, normal photochromism was obsd.,  
but the rate of decoloration was considerably smaller than that  
obsd. for the decoloration in org. solvents of similar polarity,  
indicating that the decoloration process was controlled not only by  
polarity but also by viscosity of the environment.
- IT 9002-92-0, ET 170  
(photochromic properties of spiropyran in micelle soln. of)

RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)




IT 67-66-3, Chloroform, properties

(solvent effect of, on photochromic properties of spiropyrans)

RN 67-66-3 HCAPLUS

CN Methane, trichloro- (9CI) (CA INDEX NAME)



CC 74-1 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)

ST photochromism spiropyran solvent micelle polymer film; photochromic **reaction** spiropyran substituent medium effect; spirobenzopyranindoline deriv photochromism medium polarity viscosity

IT 1119-94-4, Dodecyltrimethylammonium bromide 9002-92-0, ET 170 25155-30-0, Sodium dodecylbenzenesulfonate (photochromic properties of spiropyrans in micelle soln. of)

IT 92990-88-0P, 1,3,3-Trimethyl-5,7-dimethoxy-2-methyleneindole 121807-35-0P, 5,7-Dimethoxy-2,3,3-trimethyl-3H-indole 149987-73-5P, 1,2,3,3-Tetramethyl-5,7-dimethoxy-3H-indolium iodide (prepn. and **reaction** of, in synthesis of photochromic spiropyrans)

IT 64-17-5, Ethanol, properties 67-66-3, Chloroform, properties 123-91-1, 1,4-Dioxane, properties (solvent effect of, on photochromic properties of spiropyrans)

L71 ANSWER 16 OF 24 HCAPLUS COPYRIGHT 2002 ACS

1992:565381 Document No. 117:165381 Response of the ke test to NCI/NTP-screened chemicals. II. Genotoxic carcinogens and non-genotoxic non-carcinogens. Bakale, George; McCreary, Richard D. (Dep. Radiol., Case West. Reserve Univ., Cleveland, OH, 44106-5000, USA). Carcinogenesis, 13(8), 1437-45 (English) 1992. CODEN: CRNGDP. ISSN: 0143-3334.

AB A physicochem. carcinogen-screening test was used to measure the rate consts. of electron attachment, kes, of 105 chems. that had been screened in long-term rodent bioassays and short-term in vitro tests by the NCI/NTP. In the ke test, a pulse-cond. technique is used to generate and monitor the decay of excess electrons that serve as nucleophilic surrogates for the target tissue of rodents.

Of the 61 chems. that had been found to be rodent carcinogens as well as Salmonella mutagens, 36 yield kes that are equal to or greater than the diffusion-controlled ke of carbon tetrachloride and are considered to be pos. ke test responses. In contrast, 29 of the remaining 44 chems. that are putative non-carcinogens and non-mutagens yield kes that are neg. ke test responses. These results are combined with the ke responses of 46 non-mutagenic carcinogens and 20 mutagenic non-carcinogens that were reported earlier and are evaluated to det. the degree to which the measure of electron-accepting capacity that ke provides complements or overlaps the electrophilicity or DNA **reactivity** of chems. that is indicated by pos. mutagenicity responses in the Ames Salmonella tester strains or by pos. structural alerts, S/As, of the chems. The combined ke test results indicate that the overall predictivity of the ke test is comparable to and complements the Ames Salmonella test and S/As in identifying rodent carcinogens. Moreover, the electrons serve as non-discriminate nucleophilic targets for both genotoxic and non-genotoxic electron-accepting mols. and appear to attach with equal efficiency to carcinogens that are active in various tissues of rodents. This property of excess electrons suggests that the predictivity of the ke test could be enhanced by combining the measured ke with an appropriate lipophilicity or pharmacokinetic parameter. A prechem. electron-transfer step that had been proposed to precede chem. interactions between the carcinogen and target tissue is discussed in light of recent developments in electron-donor/-acceptor chem. and in the application of structure-activity relationships to identify carcinogens.


IT 58-89-9, Lindane 9002-92-0

(electron attachment rate const. of, genotoxic and nongenotoxic carcinogen screening in relation to)

RN 58-89-9 HCAPLUS


CN Cyclohexane, 1,2,3,4,5,6-hexachloro-, (1.alpha.,2.alpha.,3.beta.,4.alpha.,5.alpha.,6.beta.)- (9CI) (CA INDEX NAME)

Relative stereochemistry.



RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



CC 4-1 (Toxicology)

IT 51-03-6, Piperonyl butoxide 56-72-4, Coumaphos 58-89-9,  
 Lindane 59-96-1, Phenoxybenzamine 62-73-7, Dichlorvos 64-77-7,  
 Tolbutamide 69-65-8, D-Mannitol 72-20-8, Endrin 72-43-5,  
 Methoxychlor 75-35-4, Vinylidene chloride, biological studies  
 76-87-9, Triphenyltin hydroxide 77-79-2, 3-Sulfolene 78-87-5,  
 1,2-Dichloropropane 82-28-0, 1-Amino-2-methylantraquinone  
 82-68-8, Pentachloronitrobenzene 85-44-9, 1,3-Isobenzofurandione  
 88-96-0, Phthalamide 89-25-8 90-04-0, o-Anisidine 90-41-5,  
 2-Aminobiphenyl 90-94-8, Michler's ketone 91-93-0 94-20-2,  
 Chlorpropamide 94-52-0, 6-Nitrobenzimidazole 95-06-7, Sulfallate  
 95-50-1, o-Dichlorobenzene 95-53-4, biological studies 95-74-9,  
 3-Chloro-p-toluidine 95-80-7, 2,4-Diaminotoluene 95-83-0,  
 4-Chloro-o-phenylenediamine 96-12-8, 1,2-Dibromo-3-chloropropane  
 97-77-8 99-55-8, 5-Nitro-o-toluidine 99-59-2,  
 5-Nitro-o-anisidine 101-05-3, Anilazine 101-54-2,  
 N-Phenyl-p-phenylenediamine 101-77-9 101-80-4 103-33-3,  
 Azobenzene 103-85-5, 1-Phenyl-2-thiourea 105-11-3, p-Quinone  
 dioxime 105-60-2, biological studies 105-87-3, Geranyl acetate  
 106-93-4 107-06-2, 1,2-Dichloroethane, biological studies  
 108-60-1, Bis(2-chloro-1-methylethyl)ether 108-95-2, Phenol,  
 biological studies 114-86-3, Phenformin 116-06-3, Aldicarb  
 117-79-3, 2-Aminoanthraquinone 118-92-3 119-34-6,  
 4-Amino-2-nitrophenol 119-53-9, Benzoin 120-71-8, p-Cresidine  
 121-14-2, 2,4-Dinitrotoluene 121-66-4, 2-Amino-5-nitrothiazole  
 121-75-5, Malathion 122-66-7, Hydrazobenzene 124-48-1,  
 Chlorodibromomethane 126-72-7, Tris(2,3-dibromopropylphosphate)  
 127-69-5, Sulfisoxazole 128-37-0, Butylated hydroxytoluene,  
 biological studies 129-15-7, 2-Methyl-1-nitroanthraquinone  
 132-32-1, 3-Amino-9-ethylcarbazole 135-20-6, Cupferron 139-65-1,  
 4,4'-Thiodianiline 139-94-6, Nithiazide 156-10-5,  
 p-Nitrosodiphenylamine 299-42-3 303-34-4, Lasiocarpine  
 333-41-5, Diazinon 434-13-9 536-33-4, Ethionamide 542-75-6,  
 1,3-Dichloropropene 584-84-9 602-87-9, 5-Nitroacenaphthene  
 609-20-1, 2,6-Dichloro-p-phenylenediamine 842-07-9, Solvent yellow  
 14 868-85-9, Dimethyl hydrogenphosphite 968-81-0, Acetohexamide  
 999-81-5, 2-Chloroethyltrimethyl ammonium chloride 1212-29-9,  
 N,N'-Dicyclohexylthiourea 1582-09-8, Trifluralin 1634-78-2,  
 Malaoxon 1777-84-0 1836-75-5, Nitrofen 1936-15-8, Acid orange  
 10 1955-45-9, Pivalolactone 2243-62-1, 1,5-Naphthalenediamine  
 2438-88-2, 2,3,5,6-Tetrachloro-4-nitroanisole 2602-46-2, Direct  
 blue 6 2784-94-3 2832-40-8, Disperse yellow 3 3567-69-9, Acid  
 red 14 4377-33-7, 2-Chloromethylpyridine 5131-60-2,  
 4-Chloro-m-phenylenediamine 5307-14-2, 2-Nitro-p-phenylenediamine  
 8001-35-2, Toxaphene 9002-92-0 15356-70-4 17026-81-2,

3-Amino-4-ethoxyacetanilide 21739-91-3, Cytembena 62340-22-1,  
Disperse blue  
(electron attachment rate const. of, genotoxic and nongenotoxic  
carcinogen screening in relation to)

L71 ANSWER 17 OF 24 HCAPLUS COPYRIGHT 2002 ACS

1992:400277 Document No. 117:277 Mechanism of allergic cross-  
**reactions**. I. Multispecific binding of ligands to a mouse  
monoclonal anti-DNP IgE antibody. Varga, Janos M.; Kalchschmid,  
Gertrud; Klein, Georg F.; Fritsch, Peter (Dep. Dermatol., Univ.  
Innsbruck, Innsbruck, 6020, Austria). Molecular Immunology, 28(6),  
641-54 (English) 1991. CODEN: MOIMD5. ISSN: 0161-5890.


AB A recently developed solid-phase binding assay was used to  
investigate the specificity of ligand binding to a mouse monoclonal  
anti-dinitrophenyl IgE (I). All DNP-amino acids, that were tested  
inhibited the binding of the radio-labeled I to DNP covalently  
attached to polystyrene microplates; however, the concn. for 50%  
inhibition varied within four orders of magnitude, DNP-L-serine  
being the most and DNP-L-proline the least potent inhibitor. In  
addn. to DNP analogs, a large no. of drugs and other compds. were  
tested for their ability to compete with DNP for the binding site of  
I. At the concn. used for screening, 59% of compds. had no  
significant inhibition; 19% inhibited the binding of I more than  
50%. Several families of compds. (tetracyclines, polymyxins,  
phenothiazines, salicylates, and quinones) that were effective  
competitors were found. Within these families, changes in the  
functional groups attached to the family stem had major effects on  
the affinity of ligand binding. The occurrence frequencies of  
interactions of ligands with I is in good agreement with the  
semi-empirical model for multispecific antibody-ligand interactions.

IT 50-29-3, DDT, biological studies 72-54-8  
2810-69-7 9002-92-0 23488-38-2  
39568-70-2

(binding of, to anti-dinitrophenol monoclonal antibody, allergic  
cross-**reaction** mechanism in relation to)


RN 50-29-3 HCAPLUS

CN Benzene, 1,1'-(2,2,2-trichloroethylidene)bis[4-chloro- (9CI) (CA  
INDEX NAME)




RN 72-54-8 HCAPLUS

CN Benzene, 1,1'-(2,2-dichloroethylidene)bis[4-chloro- (9CI) (CA INDEX  
NAME)




RN 2810-69-7 HCAPLUS

CN Benzene, 1,2,3,4-tetrabromo-5,6-dimethyl- (9CI) (CA INDEX NAME)




RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)




RN 23488-38-2 HCAPLUS

CN Benzene, 1,2,4,5-tetrabromo-3,6-dimethyl- (9CI) (CA INDEX NAME)




RN 39568-70-2 HCAPLUS

CN Benzene, 1,2,3,5-tetrabromo-4,6-dimethyl- (9CI) (CA INDEX NAME)



IT 309-00-2, Aldrin  
 (binding of, to anti-dinitrophenol monoclonal antibody, allergic cross-**reaction** mechanisms in relation to)  
 RN 309-00-2 HCAPLUS  
 CN 1,4:5,8-Dimethanonaphthalene, 1,2,3,4,10,10-hexachloro-1,4,4a,5,8,8a-hexahydro-, (1R,4S,4aS,5S,8R,8aR)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.



CC 1-3 (Pharmacology)  
 Section cross-reference(s): 4, 15  
 ST antibody drug interaction cross **reaction**; allergy immediate hypersensitivity antibody ligand; IgE ligand cross **reaction**  
 IT Antibiotics  
 Chemicals  
 Pharmaceuticals  
 Ligands  
 Quinones  
 Sulfonamides  
 (antibodies to dinitrophenol binding by, allergic cross-**reaction** mechanism in relation to)  
 IT Hemoglobins  
 Keratins  
 Saponins  
 (binding of, to anti-dinitrophenol monoclonal antibody, allergic cross-**reaction** mechanism in relation to)  
 IT Immunoglobulins  
 (E, monoclonal, to dinitrophenol, drugs and other chem. binding to, allergic cross-**reaction** mechanism in relation to)

- IT Molecular structure-biological activity relationship  
(antibody-binding, allergic cross-reaction, of drugs  
and other chems.)
- IT Amino acids, biological studies  
(dinitrophenyl, antibodies to dinitrophenol binding by, allergic  
cross-reaction mechanism in relation to)
- IT 60-54-8D, derivs. 69-72-7D, Salicylic acid, derivs. 92-84-2D,  
Phenothiazine, derivs. 1406-11-7D, Polymyxin, derivs.  
(antibodies to dinitrophenol binding by, allergic cross-  
reaction mechanism in relation to)
- IT 50-02-2 50-10-2 50-12-4, Mesantoin 50-18-0, Endoxan  
50-29-3, DDT, biological studies 50-33-9, Phebuzine,  
biological studies 50-34-0 50-44-2, Mercaptopurine 50-65-7  
50-70-4; Sorbit, biological studies 51-05-8, Procaine  
hydrochloride 51-18-3, Tretamin 51-21-8, 5-Fluoro-uracil  
51-41-2, Levarterenol 51-67-2 51-83-2 52-01-7, Spironolactone  
52-49-3 52-62-0 52-86-8, Haloperidol 53-43-0 53-60-1,  
Promazine hydrochloride 53-86-1 54-03-5, Ustimon 54-31-9,  
Furosemid 54-32-0, Opilon 54-36-4, Metyrapone 54-42-2,  
Idoxuridine 54-49-9 54-64-8, Thimerosal 54-71-7, Pilocarpine  
hydrochloride 54-85-3 54-91-1, Pipobroman 54-92-2, Iproniazide  
54-96-6, 3,4-Pyridinediamine 55-03-8, Sodium thyroxine 55-06-1  
55-10-7 55-16-3, Scopolamine hydrochloride 55-22-1, Isonicotinic  
acid, biological studies 55-55-0, Metol 55-65-2 55-92-5,  
Methacholine 55-97-0 56-38-2, Parathion 56-55-3,  
Benz[a]anthracene 56-92-8 57-48-7, Levulose, biological studies  
57-50-1, Saccharose, biological studies 57-53-4 57-64-7  
57-66-9, Probenecid 57-83-0, Proluton, biological studies  
57-87-4 57-94-3 58-14-0, Pyrimethamine 58-15-1, Pyramidon  
58-20-8, Testosterone-cyclopentylpropionate 58-27-5, Menadione  
58-28-6, Desipramine hydrochloride 58-32-2, Persantin 58-38-8,  
Prochlorperazine 58-39-9, Perphenazine 58-46-8 58-54-8  
58-55-9, Theophyllin, biological studies 58-73-1, Diphenhydramine  
58-74-2 58-86-6, D(+)-Xylose, biological studies 59-05-2,  
Methotrexate 59-06-3, Ethopabate 59-32-5, Synopen 59-33-6,  
Pyrilamine maleate 59-43-8, Thiamine, biological studies 59-47-2  
59-63-2, Marplan 59-66-5, Diamox 59-97-2 59-99-4, Prostigmine  
60-56-0, Methimazole 60-57-1, Dieldrin 60-80-0 60-81-1  
60-89-9, Pacatal 60-99-1, Levomepromazine 61-12-1 61-16-5  
61-25-6, Papaverine hydrochloride 61-68-7 61-96-1, Nordefrin  
hydrochloride 62-13-5 62-44-2, Phenacetine 62-55-5,  
Thioacetamide 62-59-9, Veratrine 62-68-0, Proadifen  
hydrochloride 63-42-3, Lactose 63-45-6, Primaquine phosphate  
63-74-1, Sulfanilamide 63-92-3, Phenoxybenzamine hydrochloride  
64-10-8, Phenylurea 64-47-1, Physostigmine sulfate 64-55-1,  
Mebutamat 64-77-7, Tolbutamide 65-19-0 65-28-1 65-29-2  
65-45-2, Salicylamide 65-49-6, PAS 65-85-0, Benzoic acid,  
biological studies 65-86-1, Orotic acid 66-22-8, Uracil,  
biological studies 66-76-2 67-45-8, Furoxon 67-92-5,  
Dicycloverin hydrochloride 67-96-9 68-11-1, Thioglycolic acid,  
biological studies 68-91-7 68-94-0, Hypoxanthine 69-14-7,  
Trimanyl 69-65-8, Mannit 69-72-7, Salicylic acid, biological



studies 69-79-4, Maltose 69-89-6, Xanthine 69-96-5 70-30-4  
 71-82-9 72-14-0, Sulfathiazol 72-54-8 72-80-0,  
 Sterosan 76-09-5 76-20-0 76-57-3 76-61-9, Thymolblue  
 76-62-0 76-78-8 76-83-5 76-84-6, Triphenylcarbinol 77-03-2,  
 Sedulon 77-04-3, 3,3-Diethyl-2,4-dioxotetrahydropyridine 77-09-8  
 77-21-4, Doriden 77-41-8 77-67-8 78-11-5 78-28-4 79-55-0  
 79-93-6, Phenaglycodol 80-49-9 80-69-3, Tartronic acid 80-71-7  
 80-97-7 81-07-2, Saccharine 81-23-2, Dehydrocholic acid  
 81-54-9 81-60-7 81-61-8 81-89-0 81-92-5, Egmol 82-66-6,  
 Diphenadione 82-71-3 83-12-5, 2-Phenyl-1,3-indandione 83-34-1,  
 3-Methylindole 83-44-3, Desoxycholic acid 83-46-5 83-48-7  
 83-63-6, Pellidol 83-70-5, Vitamin K5 83-73-8,  
 Diiodohydroxyquinoline 83-75-0, Euquinine 83-87-4 83-88-5,  
 Lactoflavine, biological studies 84-02-6 84-11-7,  
 Phenanthraquinone 84-12-8, Entobex 84-88-8, 8-Hydroxyquinoline-5-  
 sulfonic acid 85-01-8, Phenanthrene, biological studies 86-30-6,  
 N-Nitrosodiphenylamine 86-34-0 86-48-6, 1-Hydroxy-2-naphthoic  
 acid 86-55-5, 1-Naphthalenecarboxylic acid 86-73-7, Fluorene  
 86-78-2 86-86-2, 1-Naphthylacetamide 87-41-2, Phthalide  
 87-51-4, 1H-Indole-3-acetic acid, biological studies 87-52-5  
 87-66-1, Pyrogallol 87-79-6, Sorbose 87-89-8, Inosite 88-20-0  
 88-67-5 88-96-0, Phthalamide 88-99-3, 1,2-Benzenedicarboxylic  
 acid, biological studies 89-05-4, Pyromellitic acid 89-25-8  
 89-51-0, Homophthalic acid 89-68-9, Chlorthymol 89-86-1  
 90-01-7, Saligenin 90-05-1 90-15-3, 1-Naphthalenol 90-30-2  
 90-33-5, 4-Methyl-umbelliferone 90-43-7, [1,1'-Biphenyl]-2-ol  
 90-46-0, Xanthidrol 90-47-1, Xanthone 90-50-6,  
 3,4,5-Trimethoxycinnamic acid 90-64-2, Amygdalic acid 90-82-4  
 90-85-7, Benzylephedrine 90-94-8 91-01-0, Diphenylcarbinol  
 91-10-1 91-15-6, 1,2-Benzenedicarbonitrile 91-20-3, Naphthalene,  
 biological studies 91-40-7 91-56-5, Isatin 91-57-6 91-60-1,  
 2-Naphthalenethiol 91-80-5, Methapyrilene 91-89-4, Paracotoin  
 92-24-0, Naphthacene 92-44-4, 2,3-Naphthalenediol 92-62-6,  
 Proflavine 92-69-3, [1,1'-Biphenyl]-4-ol 92-71-7, PPO 92-77-3  
 92-88-6, [1,1'-Biphenyl]-4,4'-diol 93-04-9 93-09-4,  
 2-Naphthalenecarboxylic acid 93-14-1 93-18-5, Nerolin II  
 93-35-6, Umbelliferon 93-42-5, Thionalid 93-44-7 93-56-1,  
 Phenylethylene glycol 93-60-7 94-13-3, Propyl-4-hydroxy benzoate  
 94-14-4, Cycloform 94-15-5, Larocaine 94-18-8, Nipabenzyl  
 (binding of, to anti-dinitrophenol monoclonal antibody, allergic  
 cross-reaction mechanism in relation to)  
 IT 94-19-9, Globucid 94-62-2, Piperine 94-67-7 95-04-5, Ectylurea  
 95-15-8, Thionaphthene 95-20-5 95-48-7, biological studies  
 95-63-6, Pseudocumene 95-80-7 95-87-4 95-93-2, Durol 96-91-3  
 97-05-2, Sulfosalicylic acid 97-24-5 97-65-4, biological studies  
 98-71-5 98-79-3, 5-Oxo-L-proline 98-92-0, 3-Pyridinecarboxamide  
 98-96-4, Pyrazinamide 99-06-9, biological studies 99-14-9,  
 Tricarballic acid 99-20-7 99-26-3, Dermatol 99-50-3,  
 3,4-Dihydroxybenzoic acid 99-53-6, 2-Methyl-4-nitrophenol  
 99-76-3, Nipagin 99-93-4 99-96-7, biological studies 100-88-9  
 100-97-0, biological studies 101-31-5 101-38-2 101-99-5,  
 Phenylurethane 102-08-9, Diphenylthiourea 102-98-7,

Phenylmercuryborate 103-01-5, N-Phenylglycine 103-03-7,  
1-Phenylsemicarbazide 103-16-2, Monobenzon 103-41-3, Cinnamic  
acid benzylester 103-82-2, Phenylacetic acid, biological studies  
103-85-5, Phenylthiourea 103-90-2 104-15-4, biological studies  
105-67-9 106-44-5, biological studies 106-48-9 106-49-0,  
p-Toluidine, biological studies 106-50-3, 1,4-Benzenediamine,  
biological studies 107-97-1, Sarcosine 108-39-4, m-Cresole,  
biological studies 108-45-2, 1,3-Benzenediamine, biological  
studies 108-73-6, Phloroglucine 109-57-9, Thiosinamine  
110-44-1 110-85-0, Piperazine, biological studies 111-20-6,  
Decanedioic acid, biological studies 112-47-0, 1,10-Decanediol  
112-72-1, Tetradecanol 112-86-7 113-22-4, Styptanon 113-52-0  
113-59-7, Taractan 114-80-7 114-86-3, Phenformin 115-24-2,  
Sulfonal 115-33-3 115-77-5, biological studies 116-38-1,  
Edrophonium-chloride 117-10-2, Istizin 117-34-0, Diphenylacetic  
acid 117-89-5, Trifluoperazin 118-41-2, 3,4,5-Trimethoxybenzoic  
acid, biological studies 118-55-8, Salol 118-57-0, Salophen  
118-76-3 118-79-6 118-82-1 119-39-1, 1(2H)-Phthalazinone  
119-58-4 119-90-4 119-91-5, 2,2'-Biquinoline 119-93-7  
120-14-9, Veratrumaldehyde 120-18-3, 2-Naphthalenesulfonic acid  
120-22-9, p-Nitrosodiethylaniline 120-46-7, Dibenzoylmethane  
120-57-0, Heliotropine 120-72-9, 1H-Indole, biological studies  
120-78-5 120-97-8 121-57-3 121-82-4, Hexogen 122-25-8  
122-39-4, Diphenylamine, biological studies 122-59-8 122-66-7  
122-69-0, Styracin 123-31-9, Hydroquinone, biological studies  
123-47-7, Endoiodin 124-43-6, Ortizon 124-76-5, Isoborneol  
124-87-8 125-13-3 125-33-7, Primidone 125-46-2 125-51-9  
125-52-0, Oxyphencyclimine hydrochloride 125-64-4, Noludar  
125-85-9, Parpanit 125-99-5, Pathilon 126-02-3, Cycrimine  
hydrochloride 126-27-2, Oxethazaine 126-52-3, Ethinamate  
126-81-8, Dimedon 127-48-0, Tridione 128-13-2 129-00-0,  
Pyrene, biological studies 129-20-4, Tanderil 129-77-1  
130-15-4, 1,4-Naphthalenedione 130-61-0 131-01-1, Deserpidine  
131-28-2 131-49-7, Angiografin 131-73-7 132-18-3 132-53-6  
132-86-5, 1,3-Naphthalenediol 133-10-8 133-32-4,  
1H-Indole-3-butanoic acid 133-67-5, Trichlormethiazide 134-31-6  
134-71-4, Ephetonin 135-02-4 135-19-3, 2-Naphthalenol,  
biological studies 135-31-9 135-44-4 135-88-6 136-38-9  
136-40-3, Phenazopyridine hydrochloride 136-47-0 136-72-1,  
Piperic acid 136-77-6, 4-Hexylresorcine 136-82-3, Metycaine  
137-26-8, Thiram 137-58-6, Xylocaine 138-14-7, Desferal  
138-41-0, Benzoic acid-p-sulfamide 139-33-3, Komplexon III  
139-62-8 140-22-7, Diphenylcarbazine 140-64-7 141-82-2,  
Malonic acid, biological studies 142-63-2, Piperazine hexahydrate  
143-66-8 143-67-9, Vinblastine sulfate 143-92-0, Tropenzilium  
bromide 144-75-2, Diasone 146-22-5, Mogadon 146-48-5,  
Quebrachin 146-56-5 147-24-0, Diphenhydramine hydrochloride  
147-93-3, Thiosalicylic acid 147-94-4 148-24-3, 8-Quinolinol,  
biological studies 148-72-1, Pilocarpine nitrate 148-79-8,  
Thiabendazole 149-30-4, 2-Mercaptobenzthiazole 149-91-7, Gallic  
acid, biological studies 150-69-6 150-76-5,  
Hydroquinonemonomethyl ether 150-78-7 151-83-7, Methohexital

152-02-3 152-11-4, Isoptin hydrochloride 152-72-7, Sintrom  
 154-69-8, Pyribenzamine hydrochloride 155-09-9 155-41-9  
 191-48-0, Diacenaphtho[1,2-j:1',2'-l]fluoranthene 206-44-0,  
 Fluoranthene 217-59-4, Triphenylene 288-32-4, Imidazole,  
 biological studies 288-47-1, Thiazole 298-81-7, Meladinine  
 298-96-4, Triphenyl tetrazolium chloride 299-39-8 299-42-3  
 300-48-1 302-70-5, Mitomen 302-79-4, Tretinoin 303-25-3,  
 Cyclizine hydrochloride 303-69-5, Dominal 304-84-7, Etamivan  
 306-03-6 306-07-0 306-19-4 306-21-8, Paredrine hydrobromide  
 314-03-4 315-80-0, Dibenzepin hydrochloride 317-34-0 318-98-9,  
 Propranolol hydrochloride 319-89-1, Tetrahydroxyquinone 320-77-4  
 331-39-5 339-43-5 341-70-8 350-12-9, Sulbentin 357-66-4,  
 Spirilene 360-68-9, Koprosterin 364-62-5, Metoclopramide  
 365-26-4 382-67-2 389-08-2, Nalidixic acid 390-64-7, Segontin  
 396-01-0 434-13-9 435-97-2, Marcoumar 439-14-5, Valium  
 443-48-1, Metronidazole 447-05-2, Pyridoxine phosphate 452-86-8  
 456-59-7, Cycandelate 458-24-2, Fenfluramine 466-06-8,  
 Proscillaridin 466-09-1, Uzarigenin 471-47-6, Oxamidic acid  
 475-25-2, Hematine 477-93-0, Dimethoxanate 479-18-5, Isophyllen  
 479-27-6, 1,8-Naphthalenediamine 479-92-5 480-68-2 481-06-1,  
 Santonin 481-85-6, 2-Methyl-1,4-naphthohydroquinone 482-05-3,  
 Diphenic acid 482-68-8, Sarpagan-10,17-diol 482-74-6 483-84-1,  
 Flavianic acid 484-11-7, Neocuproin 484-23-1, Nepresol  
 (binding of, to anti-dinitrophenol monoclonal antibody, allergic  
 cross-reaction mechanism in relation to)  
 IT 485-34-7, Novatophan 486-25-9, Fluorenone 486-67-9, Mersalyl  
 acid 486-79-3, Dipyroctyl 487-53-6, Oxyprocaine 488-41-5  
 489-98-5, Picramide 490-79-9, 2,5-Dihydroxybenzoic acid  
 490-91-5, Thymoquinone 490-98-2, Salicaine 492-22-8,  
 9H-Thioxanthen-9-one 492-41-1 492-70-6 493-52-7, Methyl Red  
 493-80-1, Luvistin 495-69-2, Hippuric acid 497-25-6,  
 2-Oxazolidinone 497-59-6, Meconic acid 497-75-6 498-23-7,  
 Citraconic acid 498-24-8, Mesaconic acid 500-38-9 500-89-0,  
 Thiambutosin 501-52-0, Benzenepropanoic acid 501-65-5, Tolan  
 501-81-5, 3-Pyridylacetic acid 504-15-4, Orcin 505-48-6, Suberic  
 acid 505-54-4, Hexadecanedioic acid 510-74-7, Spiramide  
 511-13-7, Detigon 511-46-6, Keithon 512-69-6, Raffinose  
 515-64-0, Elkosin 515-96-8, Semioxamazide 516-95-0 517-28-2,  
 Hematoxyline 519-05-1, Opianic acid 519-32-4 519-37-9  
 519-41-5 519-73-3, Triphenyl methane 520-07-0 520-26-3,  
 Hesperidine 521-74-4 522-12-3 522-23-6, Metofenazate  
 difumarate 522-51-0 522-66-7, Hydroquinine 523-21-7 523-87-5  
 524-42-5, 1,2-Naphthoquinone 525-48-4 526-75-0 527-06-0,  
 D-glycero-D-galacto-Heptitol 528-21-2, Gallacetophenone 530-78-9  
 532-03-6, Methocarbamol 532-54-7 532-76-3, Hexylcaine  
 hydrochloride 533-06-2 533-10-8 533-63-1, Iodival 533-73-3,  
 1,2,4-Benzenetriol 534-87-2, Effortil 535-83-1 536-21-0,  
 Norfenefrine 536-33-4, Iridozin 537-12-2 537-26-8 537-45-1  
 538-02-3, Cyclopentamine hydrochloride 538-24-9, Trilaurin  
 538-32-9 538-56-7 538-58-9 538-62-5, Diphenylcarbazone  
 539-08-2 539-09-3, Elbon 539-21-9, Iversal 539-47-9  
 543-15-7, Heptaminol hydrochloride 547-57-9 547-58-0,

Methylorange 548-00-5, Tromexan 548-35-6 548-57-2, Miracil D 548-73-2, Dehydrobenzperidol 550-70-9, Triprolidine hydrochloride 550-74-3 552-22-7, Thymoliodide 552-46-5, .alpha.-Naphthylamine hydrochloride 552-94-3, Diplosal 553-06-0 553-30-0, Proflavinhemisulfate 554-24-5, Phenobutiodil 554-57-4, Methazolamide 555-28-2, Eucaïne hydrochloride 555-30-6, Methyldopa 555-68-0, m-Nitrocinnamic acid 556-50-3 561-20-6 562-09-4, Systral 562-10-7, Doxylamine 569-51-7, Hemimellitic acid 569-59-5 569-61-9, Pararosaniline 571-60-8, 1,4-Naphthalenediol 575-44-0, 1,6-Naphthalenediol 576-68-1, Mannomustin 578-19-8, Diaspirin 578-94-9 579-10-2, Exalgin 579-23-7, Cyclovalone 579-75-9, Salicylic acid methylether 580-16-5, 6-Quinolinol 581-43-1, 2,6-Naphthalenediol 582-17-2, 2,7-Naphthalenediol 588-59-0, Stilbene 590-63-6, Urecholine chloride 594-14-9 599-04-2 603-00-9 603-50-9, Dulcolax 603-52-1, Diphenylurethane 603-63-4, Hypnal 603-64-5 604-75-1, Oxazepam 606-90-6, Piprinhydrinate 607-00-1 608-59-3, Gluconate 608-66-2, Melampyrit 611-36-9, 4-Quinolinol 611-71-2 612-41-9, o-Nitrocinnamic acid 612-52-2 613-03-6 613-12-7 613-31-0, Dihydroanthracene 614-33-5 620-40-6, Tribenzylamine 620-99-5, Holocaine hydrochloride 621-06-7, Phenylacetanilide 621-08-9 621-37-4 621-71-6, Tricaprin 621-82-9, Cinnamic acid, biological studies 622-03-7, Diphenylthiocarbazine 622-64-0 626-64-2, 4-Pyridinol 632-25-7, o-Sulfobenzoic acid 635-41-6, Trimetozin 637-56-9, p-Phenetidine hydrochloride 637-58-1, Tronothane hydrochloride 644-26-8, Stovain 644-62-2 655-05-0, Thozalinone 673-31-4, Phenprobamate 693-23-2, Decamethylenedicarboxylic acid 696-62-8 697-91-6 699-06-9 712-48-1 721-19-7, Methastyridone 729-99-7 738-70-5, Trimethoprim 739-71-9, Trimipramine 742-20-1, Cyclopenthiazine 749-02-0 749-13-3, Trifluperidol 751-94-0 826-39-1, Mecamylamine hydrochloride 830-81-9, 1-Naphthylacetate 840-65-3 841-32-7, Diphenylpropylacetic acid 846-49-1, Lorazepam 846-50-4, Temazepam 848-75-9, Lormetazepam 852-19-7, Sulfapyrazole 853-34-9, Ketophenyl butazone 868-18-8 877-43-0, 2,6-Dimethylquinoline 890-98-2, Mandelic acidbenzylester 891-33-8, Tutocaine 894-71-3, Nortriptyline hydrochloride 897-15-4, Dosulepin hydrochloride 909-39-7, Insidon 910-86-1 912-60-7, Narcotine hydrochloride 933-67-5, 7-Methylindole 938-25-0, 1,2-Naphthalenediamine 942-46-1, Methylephedrine hydrochloride 943-17-9 953-26-4 956-48-9 957-51-7, Diphenamid 958-93-0, Thenyldiamine hydrochloride 963-39-3, Demoxepam 964-52-3 965-52-6, Nifuroxazide 967-80-6 969-33-5, Cyproheptadine hydrochloride 977-79-7 982-43-4 985-13-7 1007-33-6 1011-54-7, trans-2-Methoxy cinnamic acid 1011-92-3, .alpha.-Cyano-cinnamic acid 1015-89-0, 6(5H)-Phenanthridinone 1021-11-0 1059-28-5, Tussukal 1070-95-7, Guanoctine hydrochloride 1077-28-7, DL-Thioctic acid 1079-71-6, Octahydroanthracene 1082-57-1, Rhinogutt 1083-57-4 1088-92-2 1094-08-2, Profenamine hydrochloride 1098-60-8 1115-70-4, Metformin hydrochloride 1142-42-3 1145-36-4, Felogen 1146-95-8, Gilutensin hydrochloride 1147-56-4,

1-(2-Thiazolylazo)-2-naphthol 1151-11-7, Solu-Biloptin  
 1155-49-3, Falicain 1156-19-0, Tolazamide 1172-18-5, Flurazepam  
 dihydrochloride 1173-88-2, Stapenor 1176-03-0, Psicaine  
 1176-08-5, Phenyltoloxamine citrate 1179-69-7, Torecan  
 1197-18-8, Tranexamic acid 1198-55-6 1199-77-5,  
 .alpha.-Methyl-cinnamic acid 1211-28-5, Katovit 1212-72-2,  
 Mephen-termine sulfate 1215-83-4, Silomat 1218-35-5 1225-55-4  
 1228-19-9, Glypinamide

(binding of, to anti-dinitrophenol monoclonal antibody, allergic  
 cross-reaction mechanism in relation to)

IT 1229-29-4, Doxepin hydrochloride 1229-69-2 1240-15-9,  
 Propiomazine hydrochloride 1248-42-6 1256-01-5, Pasaden  
 1323-64-4 1391-57-7, Filmaron 1400-62-0, Orcein 1407-14-3,  
 Helenine 1421-86-9, Strychnine hydrochloride 1480-19-9,  
 Fluanisone 1491-41-4, Maretin 1508-27-6 1508-75-4 1508-76-5,  
 Procyclidine hydrochloride 1582-09-8, Trifluraline 1596-70-9  
 1622-62-4, Flunitrazepam 1622-79-3 1639-60-7 1641-74-3,  
 Nicametate citrate 1642-54-2 1674-48-2 1701-77-5,  
 Methoxy-phenylacetic acid 1722-62-9 1740-22-3, Surexin  
 1764-85-8, Epithiazide 1786-81-8, Prilocaine hydrochloride  
 1837-57-6, Rivanol 1841-19-6, Fluspirilene 1847-63-8, Nafoxidine  
 hydrochloride 1867-66-9, Ketamine hydrochloride 1893-33-0  
 1944-12-3, Fenoterol hydrobromide 1975-50-4, 2-Methyl-3-nitro-  
 benzoic acid 2002-29-1 2043-43-8, Lactamide 2053-26-1,  
 Papaverine sulfate 2062-78-4 2081-65-4 2135-17-3, Flumethason  
 2139-47-1 2181-22-8 2192-20-3 2210-63-1, Monophenylbutazone  
 2217-44-9 2218-94-2, Nitron 2243-62-1, 1,5-Naphthalenediamine  
 2259-96-3, Cyclothiazide 2295-31-0, 2,4-Thiazolidinedione  
 2307-81-5, Quiloflex 2313-87-3, Ethoxazene hydrochloride  
 2315-02-8 2321-07-5 2324-94-9 2347-80-0 2348-17-6  
 2350-32-5, Stadacaine 2398-96-1, Tonofal 2435-53-2 2444-46-4  
 2447-57-6, Fanasil 2448-68-2 2451-01-6, Terpinhydrate  
 2465-59-0, Oxypurinol 2525-09-9 2530-97-4 2572-61-4  
 2574-78-9 2606-93-1 2608-24-4 2622-26-6, Periciazine  
 2691-46-5, Spasmo-Papamid 2709-56-0, Flupenthixol 2773-92-4,  
 Quotane hydrochloride 2784-55-6 2810-69-7 2825-60-7,  
 Formocortal 2870-71-5, Tropin 2898-12-6, Medazepam 2955-38-6,  
 Prazepam 2961-04-8 2987-16-8 3069-07-6, Tetraacetyl hydrazine  
 3093-35-4, Halcinonide 3112-31-0, 1H-Pyrazole-3,5-dicarboxylic  
 acid 3115-05-7 3254-89-5, Diphenidol hydrochloride 3269-83-8  
 3339-11-5, Tolpropamine hydrochloride 3403-42-7, Methopromazine  
 maleate 3413-58-9, Optochin hydrochloride 3416-26-0, Lidoflazine  
 3453-83-6 3458-28-4, D(+)Mannose 3459-20-9, Redul 3546-03-0  
 3546-41-6 3605-01-4, Piribedil 3614-30-0 3614-69-5  
 3615-41-6, Rhamnose 3682-32-4, 2-Nitroso-1-naphthol-4-sulfonic  
 acid 3685-84-5, Meclophenoxate hydrochloride 3688-85-5,  
 Diapamide 3689-50-7, Oxomemazin 3717-88-2 3731-59-7  
 3735-45-3, Monzal 3735-90-8, Fencarbamide 3759-07-7 3772-76-7,  
 Methofadin 3811-56-1, Surfen 3818-88-0, Tricyclamol chloride  
 3829-86-5 3871-82-7, Methylperidol hydrochloride 3937-56-2,  
 1,9-Nonanediol 3963-81-3, Trigemin 4003-94-5, 4-Nitrostilbene  
 4044-65-9, 1,4-Phenylenediisothiocyanate 4247-16-9,

Methylglutamine 4330-99-8 4394-00-7 4427-56-9, Isothymol  
 4546-48-9 4551-59-1 4582-18-7, Endomid 4682-36-4 4724-59-8  
 4839-46-7, 3,3-Dimethylglutaric acid 4991-65-5, Tioxolon  
 5034-76-4, Indoxole 5041-09-8, Isobutylamine hydrochloride  
 5076-82-4, Sarcosine anhydride 5144-52-5 5255-68-5 5322-53-2,  
 Oxiperomide 5370-01-4 5416-45-5, Phenyldiphenylcarbamate  
 5428-54-6, 2-Methyl-5-nitrophenol 5437-38-7, 3-Methyl-2-nitro-  
 benzoic acid 5449-84-3 5536-17-4 5579-13-5 5585-60-4  
 5588-29-4, Fenmetramide 5588-31-8, Imidoline hydrochloride  
 5591-29-7, Etafedrine hydrochloride 5591-45-7, Thiiothixene  
 5667-46-9, Dioxyline phosphate 5716-20-1, Vasculat 5870-29-1,  
 Cyclopentolate hydrochloride 5874-97-5 5875-06-9 5936-29-8  
 5969-39-1 5987-82-6, Novesin 6027-28-7, Hostacaine 6028-35-9  
 6038-78-4 6056-11-7, Selvigon 6114-26-7, Veritol sulfate  
 6138-47-2 6153-33-9 6164-87-0, Ronicol 6190-43-8, Helmitol  
 6192-92-3 6202-05-7, Cyclomethycaine sulfate 6202-23-9,  
 Cyclobenzaprine hydrochloride 6217-24-9 6284-40-8, Meglumin  
 6411-75-2 6452-71-7 6493-05-6, 3,7-Dimethyl-1-(5-  
 oxohexyl)xanthine 6506-37-2, Nimorazol 6521-30-8 6536-18-1  
 6556-11-2, Inositol nicotinate 6575-24-2 6673-35-4, Practolol  
 6700-56-7, Ethoheptazine citrate 6724-53-4, Perhexiline maleate  
 7008-15-3 7009-43-0 7085-55-4, Troxerutin 7125-73-7,  
 Flumetramide 7195-27-9, Mefrusid 7199-29-3, Cyheptamide  
 7210-92-6, Tolycaine hydrochloride 7242-04-8, Pentaacetyl gitoxin  
 7270-12-4, Resotren 7297-25-8 7413-36-7, Nifenalol 7414-95-1  
 7491-74-9, Piracetam 7517-19-3, L-Leucine methyl ester  
 hydrochloride 7601-55-0, Metubine iodide 8002-89-9 8006-08-4,  
 Ergotoxinine 8015-17-6, Metrotonin 8015-18-7, Veramon  
 8064-60-6, Primulin 9001-22-3, Emulsin 9002-92-0  
 9004-53-9, Dextrin 9005-79-2, Glycogen, biological studies  
 9005-80-5, Inulin 10040-45-6 10176-39-3 10238-21-8  
 10347-81-6 10402-53-6 10405-02-4 10539-19-2, Eupaverin  
 10563-70-9, Melitracen hydrochloride 10592-03-7, Vincamine  
 hydrochloride 11014-59-8, Lanatoside 11024-24-1 12002-15-2,  
 Sapamine 12041-92-8, Hexyltheobromine 12694-25-6, Bi-9H-fluorene  
 13055-82-8, Reproterol hydrochloride 13422-16-7, Triflocin  
 13472-79-2 13492-01-8 13523-86-9, Pindolol 13636-10-7  
 13636-18-5 13665-88-8, Mopidamol 13900-17-9 13977-28-1  
 14198-59-5 14222-60-7, Ektebin 14255-87-9, Parbendazole  
 14293-44-8 14516-56-4 14538-56-8, Piperazine phosphate  
 14543-76-1 14698-29-4, Oxolinic acid 14759-06-9 15307-79-6,  
 Voltaren

(binding of, to anti-dinitrophenol monoclonal antibody, allergic  
 cross-reaction mechanism in relation to)

IT 15308-34-6, Novadral hydrochloride 15351-13-0, Nicofuranose  
 15402-76-3 15537-73-2, Silubin 15585-38-3 15676-16-1,  
 Sulpiride 15687-27-1, Ibuprofen 15825-70-4 15876-67-2  
 16110-98-8, Phenyl maleic acid 16662-46-7, Gallopamil  
 hydrochloride 16773-42-5, Ornidazole 16994-56-2 17088-72-1  
 17226-75-4 17273-86-8 17297-82-4 17407-37-3,  
 DL-.alpha.-Tocopherol succinate 17509-71-6 17560-51-9,  
 Metolazone 17692-39-6 17892-25-0 18174-58-8 18559-94-9

18869-73-3, Triacetyl diphenolisatin 19188-90-0 19237-84-4,  
 Prazosin hydrochloride 19311-91-2 19387-91-8, Tinidazole  
 19562-30-2 19794-93-5, Trazodone 20153-98-4 20231-81-6, Uzarin  
 20277-92-3, N,N-Diphenyl-guanidine 20380-58-9 20423-87-4  
 20432-64-8, Iprindole hydrochloride 20455-68-9, Dibenzylamine  
 hydrochloride 20788-07-2, Terenol 20833-93-6 21361-95-5  
 21498-08-8 21535-47-7, Mianserine hydrochloride 21721-92-6  
 21738-42-1 21829-25-4, Nifedipine 22059-60-5, Disopyramide  
 phosphate 22071-15-4, Ketoprofen 22089-22-1, Trofosfamide  
 22204-53-1, Naproxen 22232-55-9, Modatrop 22254-24-6,  
 Ipratropiumbromide 22494-42-4, Diflunisal 22760-18-5 22832-87-  
 7 22888-70-6 23031-32-5, Terbutalinsulfate 23092-17-3,  
 Halazepam 23111-34-4, Feclobuzone 23142-01-0 23239-51-2  
 23288-49-5, Probutol 23307-02-0 23327-57-3, Nefopam  
 hydrochloride **23488-38-2** 23607-71-8 23694-81-7,  
 Mepindolol 23873-81-6 23983-43-9 24168-96-5 24169-02-6  
 24324-17-2, 9-Fluorenyl-methanol 24526-64-5, Nomifensin  
 24561-10-2, Piperocaine hydrochloride 24600-36-0, Fominoben  
 hydrochloride 24815-24-5 25046-79-1, Glisoxepid 25167-82-2  
 25167-84-4 25498-47-9, Saiodin 25717-80-0 25812-30-0,  
 Gemfibrozil 25953-17-7 26020-55-3 26309-95-5, Pivampicillin  
 hydrochloride 26598-44-7 26718-25-2 26864-56-2, Penfluridol  
 26921-17-5, Timolol-maleate 26983-52-8, Diphenol 27479-32-9  
 27848-84-6 28346-70-5, Naphthalenediol 28738-34-3 29094-61-9,  
 Glipizide 29110-48-3 29868-97-1 30440-92-7 30900-94-8  
 30919-08-5 31329-57-4, Naftidrofuryl 31431-39-7, Mebendazole  
 31566-31-1, Glycerin monostearate 31793-07-4, Pirprofen  
 31842-01-0, Indoprofen 31901-98-1, Naphthalenetetracarboxylic acid  
 32672-69-8 32780-64-6, Labetalol hydrochloride 32972-46-6  
 33125-97-2, Etomidate 33342-05-1 33396-37-1 33401-94-4,  
 Pyrantel-tartrate 33402-03-8 33996-33-7, Oxaceprol 34183-22-7,  
 Propafenone hydrochloride 34552-84-6, Isoxicam 34661-75-1  
 35306-33-3 35412-64-7 35604-67-2, Viloxazine hydrochloride  
 36236-67-6, Meclizine hydrochloride 36282-47-0, Tramadol  
 hydrochloride 36322-90-4, Piroxicam 36637-19-1, Etidocaine  
 hydrochloride 37275-48-2, Dipyridyl 37887-33-5 38096-29-6,  
 Pyridinediamine 38194-50-2, Sulindac 38304-91-5, Minoxidil  
 38866-78-3 39379-48-1, Reten 39461-53-5, Pyrenedione  
 39562-70-4 **39568-70-2** 40180-04-9 41100-52-1, Memantine  
 hydrochloride 41247-05-6, DL-Xylose 41340-25-4, Etodolac  
 41451-91-6, Erythromycine 41587-33-1 41767-29-7, Fluocortinbutyl  
 ester 41838-38-4 41847-77-2 41960-46-7 42200-33-9, Nadolol  
 43218-56-0 49721-50-8 49746-00-1, Twiston 50322-92-4  
 50591-64-5 50679-08-8 50832-74-1, Nifurprazine hydrochloride  
 50838-36-3 50926-65-3, Novalgin-quinine 51481-61-9 51703-77-6  
 51940-44-4, Pipemidic acid 51996-59-9 52432-72-1, Oxeladin  
 citrate 52441-07-3, Parsol 52468-60-7 52849-55-5,  
 DL-Metipranolol 53179-11-6, Loperamide 53623-34-0 53663-23-3  
 53783-83-8, Tromantadine 53859-10-2 53885-35-1, Ticlopidine  
 hydrochloride 54024-22-5 54504-70-0 54750-10-6, Isolevin  
 54767-75-8, Suloctidil 54812-66-7 54965-24-1, Tamoxifen citrate  
 55031-26-0, Iodoeosine 55268-74-1, Praziquantel 55327-22-5



56050-03-4, Meclozamine citrate 56392-17-7, Metoprolol tartrate  
 56601-85-5 57109-90-7 57808-66-9 58934-46-6, Lorcainide  
 hydrochloride 59831-65-1, Halopemide 59954-01-7 60525-15-7  
 60539-09-5 60607-34-3 61169-36-6 61229-67-2, Bromocresol red  
 63250-48-6, Piprozoline 64019-93-8 65271-80-9 65277-42-1,  
 Ketoconazole 65431-33-6, Trypaflavine 65923-65-1 66894-06-2  
 69494-65-1, Migraenin 72762-00-6, 2-Pyridinol 73548-65-9,  
 Veraethyl 74217-46-2 74347-31-2 75507-68-5 77614-18-7  
 78361-94-1, Naphthoquinoline 81098-57-9 81177-02-8 83943-60-6  
 98578-19-9 104700-83-6 109893-47-2 139352-30-0, Pergalen  
 139369-53-2 139369-54-3 139369-55-4 139369-56-5 139369-57-6  
 139369-58-7, 2H-1-Benzopyran-3,3,4,4,5,7-hexol 139369-59-8  
 139369-60-1 139369-61-2 139369-62-3 139369-64-5 139369-65-6  
 139369-66-7 139369-67-8 139369-68-9 139369-70-3 139369-71-4  
 139411-97-5 139412-02-5 139412-03-6 139412-04-7 139465-28-4,  
 Dipa-Vit B 15 139465-72-8, Nipagin T 139466-02-7, Novanal  
 139466-06-1, Phenidol 139556-82-4 139984-92-2

(binding of, to anti-dinitrophenol monoclonal antibody, allergic  
 cross-reaction mechanism in relation to)

IT 50-33-9, Butazolidine, biological studies 50-41-9, Clomiphene  
 citrate 50-48-6, Amitriptylin 50-63-5, Chloroquin diphosphate  
 50-71-5, Alloxan 50-78-2, Acetyl salicylic acid 50-85-1,  
 m-Cresotinic acid 51-17-2, Benzimidazole 51-35-4, Hydroxyproline  
 51-55-8, Atropine, biological studies 51-66-1 51-84-3,  
 biological studies 52-90-4, Cysteine, biological studies  
 54-31-9, Furosemide 54-84-2 54-95-5, Cardiazole 55-56-1,  
 Chlorhexidine 55-98-1, Busulphan 56-40-6, Glycine, biological  
 studies 56-41-7, Alanine, biological studies 56-45-1, Serine,  
 biological studies 56-55-3, Benzanthrane 56-72-4, Coumafes  
 56-75-7, Chloramphenicol 56-85-9, Glutamine, biological studies  
 56-87-1, L-Lysine, biological studies 56-89-3, Cystine, biological  
 studies 57-37-4, Benactyzine.hydrochloride 57-67-0,  
 Sulfaguanidine 57-68-1, Sulfamethazine 57-92-1, Streptomycin,  
 biological studies 58-08-2, biological studies 58-15-1  
 58-33-3, Atosil 58-61-7, Adenosine, biological studies 58-94-6,  
 Chlorthiazide 59-42-7 59-49-4, 2(3H)-Benzoxazolone 60-18-4,  
 L-Tyrosine, biological studies 60-27-5 60-32-2 60-54-8  
 60-80-0, Antipyrin 61-12-1 61-33-6, Penicillin G, biological  
 studies 61-72-3, Cloxacillin 61-75-6, Bretyliumtosylate  
 61-90-5, Leucine, biological studies 61-94-9 61-96-1, Corbasil  
 63-68-3, Methionine, biological studies 63-91-2, Phenylalanine,  
 biological studies 64-77-7, Artosin 64-86-8, Colchicine  
 64-95-9, Adiphenin 65-45-2, Salicylic acid amide 65-82-7,  
 N-Acetylmethionine 66-81-9 67-03-8, Aneurine hydrochloride  
 67-52-7, 2,4,6(1H,3H,5H)-Pyrimidinetrione 68-35-9, Sulfadiazin  
 68-41-7, D-Cycloserine 69-09-0, Chlorpromazine hydrochloride  
 69-27-2 69-53-4, Ampicillin 70-47-3, Asparagine, biological  
 studies 71-00-1, Histidine, biological studies 72-18-4, Valine,  
 biological studies 72-19-5, Threonine, biological studies  
 72-48-0, Alizarin 73-22-3, Tryptophane, biological studies  
 73-24-5, Adenine, biological studies 73-32-5, Isoleucine,  
 biological studies 73-48-3, Benzylrodiuran 74-79-3, L-Arginine,



biological studies 76-22-2 76-29-9 76-60-8, Bromcresolgreen  
76-65-3, Amolanone 77-02-1, Allonal 77-36-1 77-46-3,  
Acedapsone 77-65-6, Adalin 77-91-8 79-05-0, Propanamide  
79-57-2, Oxytetracycline 80-03-5 80-05-7, biological studies  
80-32-0, Vetisulid 80-77-3, Chlormezanone '80-97-7 81-61-8  
81-64-1, Quinizarine 82-05-3, Benzanthrone 82-45-1,  
1-Amino-anthraquinone 82-54-2, Cotarnine 82-86-0,  
1,2-Acenaphthylenedione 82-93-9 83-32-9, Acenaphthene 83-40-9,  
o-Cresotinic acid 84-65-1, Anthraquinone 85-18-7 85-73-4,  
Taleudron 86-42-0 86-54-4 86-74-8, Carbazole 87-08-1  
87-32-1, N-Acetyl-DL-tryptophan 87-88-7 88-21-1 89-56-5,  
p-Cresotinic acid 90-64-2, Amygdalic acid 90-84-6 91-33-8  
91-64-5, 2H-1-Benzopyran-2-one 93-08-3 93-10-7, Quinaldinic acid  
93-18-5 93-44-7 94-09-7, Benzocaine 94-12-2 94-19-9  
94-20-2, Chlorpropamide 94-25-7, Butesine 94-41-7 95-25-0,  
Chlorzoxazone 95-55-6, o-Aminophenol 95-85-2 96-83-3, Iopanoic  
acid 97-59-6, Allantoin 98-37-3 98-50-0, Arsanilic acid  
99-32-1, Chelidonic acid 99-91-2 101-71-3 102-07-8,  
Carbanilide 103-12-8, Prontosil 103-32-2, Benzylaniline  
103-41-3 103-84-4, N-Acetylaniline 103-89-9 103-90-2  
104-06-3, Conteben 105-20-4, Betazole 106-34-3, Quinhydrone  
106-51-4, p-Benzoquinone, biological studies 113-92-8,  
Chlorpheniramine maleate 114-83-0 115-39-9, Brom Phenol Blue  
115-51-5, Ambutoniumbromide 115-68-4, Irgamid 118-10-5  
118-23-0, Ambodryl 118-75-2, Chloranil, biological studies  
118-92-3, o-Aminobenzoic acid 119-53-9, Benzoin 119-61-9,  
Benzophenone, biological studies 120-32-1 120-34-3, Irgafen  
121-25-5, Amprolium 122-11-2, Madribon 122-80-5 124-04-9,  
Hexanedioic acid, biological studies 125-60-0, Baralgin-Amid  
126-07-8 126-43-2 127-69-5, Sulfafurazol 127-79-7,  
Sulfamerazine 127-81-1, Salthion 130-16-5 130-22-3 131-08-8  
131-09-9 132-98-9, Isocillin 134-50-9, 9-Aminoacridine  
hydrochloride 134-81-6, Benzil 134-85-0 136-95-8,  
2-Benzothiazolamine 137-08-6, Calcium-D-pantothenate 138-39-6,  
Mafenide 139-85-5 141-82-2, Propanedioic acid, biological  
studies 143-37-3, Ethanimidamide 144-83-2 145-94-8,  
Chlorindanol 146-56-5 147-24-0, Benadryl 147-55-7,  
Pheneticillin 147-85-3, Proline, biological studies 148-64-1  
148-82-3, Melfhalan 149-15-5, Butyn Sulfate 149-64-4 150-13-0,  
p-Aminobenzoic acid 150-69-6, p-Ethoxyphenylurea 152-47-6  
153-61-7, Cefalotin 154-87-0, Cocarboxylase 154-97-2,  
Contrathion 208-96-8, Acenaphthylene 218-01-9, Chrysene  
243-42-5, Benzo[b]naphtho[2,3-d]furan 260-94-6, Acridine  
298-46-4, 5H-Dibenz[b,f]azepine-5-carboxamide 298-57-7,  
Cinnarizine 303-81-1, Novobiocin 305-03-3, Chlorambucil  
309-00-2, Aldrin 313-67-7, Aristolochic acid 315-30-0,  
Allopurinol 316-41-6, Berberinsulfate 334-48-5, Decanoic acid  
346-18-9, Polythiazide 370-81-0 437-74-1 441-38-3 446-86-6,  
Azathioprine 464-41-5 466-49-9, Aspidospermin 467-22-1,  
Carbiphen hydrochloride 474-25-9 481-74-3 488-82-4,  
D-Arabinitol 489-49-6, Cetrarin 497-76-7, Arbutin 498-23-7,  
Citraconic acid 499-12-7, Aconitic acid 515-64-0, Sulfisomidine

519-87-9, Acetyldiphenylamine 520-77-4, Ethadione 521-74-4,  
 Broxyquinoline 526-08-9, Sulfaphenazole 529-65-7 530-43-8,  
 Chloromycetin palmitate 531-75-9, Aesculin 537-05-3, Acoïn  
 537-92-8 543-24-8, N-Acetylglycine 545-93-7 547-44-4  
 550-81-2, Amopyroquin

(binding of, to anti-dinitrophenol monoclonal antibody, allergic  
 cross-reaction mechanisms in relation to)

IT 551-27-9, Propicillin 555-96-4 556-08-1, p-Acetaminobenzoic acid  
 564-25-0 569-84-6 575-36-0 577-33-3, Anthrarobin 578-66-5,  
 8-Aminoquinoline 581-97-5 588-68-1 590-46-5,  
 Betaine.hydrochloride 591-07-1, N-Acetylurea 591-08-2,  
 N-Acetylthiourea 599-79-1 599-88-2, Sulfaperin 602-09-5,  
 [1,1'-Binaphthalene]-2,2'-diol 606-04-2 613-78-5, Betol  
 616-91-1, N-Acetyl-L-cysteine 625-53-6, N-Ethylthiourea  
 630-55-7, Amphotropin 636-54-4, Clopamide 637-32-1 637-49-0  
 651-06-9, Bayrena 653-03-2, Butaperazine 655-35-6 671-95-4  
 703-80-0, 3-Acetylinole 723-46-6 729-99-7 738-66-9,  
 Bis(4-nitrophenyl) carbodiimide 751-97-3, Reverin 768-90-1  
 768-94-5, Tricyclo[3.3.1.1<sup>3,7</sup>]decan-1-amine 776-34-1 776-75-0,  
 Benzoylpiperidine 795-13-1, Formyl sulfamethin 804-63-7  
 828-51-3 846-50-4 849-55-8, Buphenin hydrochloride 873-76-7  
 882-09-7 914-00-1, Methacycline 935-56-8 956-04-7 963-07-5  
 968-81-0, Acetohexamide 980-71-2 982-24-1, Clopenthixol  
 992-21-2, Lymecycline 1050-48-2, Benzilium bromide 1071-93-8  
 1072-71-5, 2,5-Dimercapto-1,3,4-thiadiazole 1098-50-6,  
 Dansylvaline 1100-21-6, Dansylisoleucine 1100-23-8 1101-68-4,  
 Dansylglutamic acid 1104-36-5, Dansylphenylalanine 1111-06-4  
 1126-46-1 1126-81-4 1134-47-0, Baclofen 1143-38-0, Cignolin  
 1145-80-8, N-Carbobenzoxo-L-serine 1163-36-6, Allercur  
 1188-01-8, DL-Alanyl-glycine 1190-53-0 1215-83-4, Clobutinol  
 hydrochloride 1218-34-4, N-Acetyl-L-tryptophan 1225-60-1,  
 Andantol 1239-94-7, Dansylproline 1248-42-6, Baralgin-Ketone  
 1249-84-9, Azacosterol.hydrochloride 1260-17-9 1263-03-2  
 1333-08-0 1397-89-3, Amphotericin B 1398-20-5, Baptisin  
 1400-61-9, Mycostatin 1401-79-2, Vionactan 1404-26-8, Polymyxin  
 B 1404-88-2, Tyrothricin 1405-97-6, Gramicidin 1477-19-6  
 1538-09-6 1622-61-3, Clonazepam 1642-81-5 1649-18-9, Azaperone  
 1684-42-0, Acranil 1798-50-1, Azacyclonol.hydrochloride  
 1812-30-2 1824-50-6 1824-52-8 1824-58-4 1861-40-1, Benefin  
 1867-58-9 1915-83-9 1951-25-3 1986-53-4 2016-88-8,  
 Amilorid.hydrochloride 2022-85-7 2043-38-1, Butizid 2058-52-8,  
 Clothiapine 2109-73-1, Butacetin 2134-24-9 2138-22-9  
 2179-37-5 2350-32-5 2432-99-7 2465-59-0, Alloxanthine  
 2483-49-0 2508-72-7 2622-30-2, Carphenazine 2667-89-2,  
 Bisbentiamine 2751-09-9, Oleandomycin triacetate 3073-87-8,  
 Dimethyl-Popop 3082-75-5 3274-19-9, 1-Acetylaminoanthraquinone  
 3287-99-8, Benzylamine hydrochloride 3347-56-6 3483-82-7  
 3485-62-9 3505-38-2, Carbinoxamine maleate 3521-62-8 3533-97-9  
 3562-84-3 3624-96-2, Bialamicol hydrochloride 3679-64-9  
 3684-46-6 3686-68-8, Cornecaine 3876-10-6 3922-90-5,  
 Oleandomycin 4205-90-7, Clonidin 4330-99-8 4361-81-3  
 4388-82-3, Barbexaclon 4393-72-0 4419-39-0 4845-99-2

4942-47-6, Tricyclo[3.3.1.1<sup>3</sup>,7]decane-1-acetic acid 4985-15-3,  
 Agedal 5068-28-0 5205-82-3, Bevoniummethysulfate 5251-34-3,  
 Cloprednol 5355-48-6 5490-27-7 5568-90-1,  
 Isopromethazine.hydrochloride 5585-73-9, Butriptyline  
 hydrochloride 5588-20-5 5591-49-1, Anilamate 5667-98-1,  
 Baludon 5786-21-0, Clozapin 5843-82-3 5892-41-1 6011-39-8  
 6469-93-8, Chlorprothixene hydrochloride 6804-07-5, Carbadox  
 6933-90-0 7421-40-1 7542-37-2, Paromomycin 7683-59-2, Aludrin  
 8000-95-1, Caffeine sodium-benzoate 8002-85-5 8044-71-1,  
 Cetrimide 8048-52-0 8059-24-3, Adermine 8068-28-8,  
 Colistimethate sodium 10199-89-0 10268-71-0 10323-20-3,  
 D(-)-Arabinose 12041-72-4, Formo-Cibazol 12246-80-9 13068-66-1  
 13296-94-1 13461-01-3 13539-59-8 13655-52-2, Alprenolol  
 13726-85-7 13838-08-9 13900-12-4 13957-27-2 14205-39-1  
 14252-80-3 14286-84-1 14358-44-2 14414-68-7, Benzidine,  
 hydrochloride 14548-46-0, 4-Benzoyl-pyridine 14976-57-9  
 15148-80-8, Bupranolol hydrochloride 15301-40-3 15318-45-3  
 15351-05-0 15686-71-2, Cefalexin 15761-39-4 16862-11-6  
 17039-58-6, Dansylmethionine 17321-77-6 17344-99-9 17671-51-1  
 17683-09-9 18109-80-3 18109-81-4 18323-44-9, Sobelin  
 18500-45-3 19461-29-1 20191-75-7 20448-86-6 21352-09-0  
 21593-23-7 21898-19-1, Clenbuterol.hydrochloride 22131-79-9  
 22198-72-7 22199-46-8 22316-47-8, Clobazam 22457-89-2  
 23067-13-2 23110-15-8 23828-92-4, Ambroxol.hydrochloride  
 23964-57-0 25122-57-0, Clobetasone butyrate 25161-41-5,  
 Acevaltrate 27043-22-7, Ethoxybenzamide 27195-22-8 27323-18-8  
 27941-88-4 28217-22-3 28217-24-5 28805-76-7, Aminobutyric acid  
 28860-95-9 28981-97-7 29022-11-5 29122-68-7, Atenolol  
 29546-59-6 30408-30-1, Nybomycin 30578-37-1 30964-13-7  
 31135-62-3, Quinolinamine 31677-93-7 32442-99-2

(binding of, to anti-dinitrophenol monoclonal antibody, allergic  
 cross-reaction mechanisms in relation to)

IT 32795-47-4, Alival 32862-97-8 32988-50-4, Viomycin 33386-08-2,  
 Buspirone hydrochloride 33817-20-8, Pivampicillin 35021-10-4  
 35021-12-6, Dansylserine 35021-16-0, Dansylthreonine 35026-16-5,  
 Dansylhydroxyproline 35408-57-2 35661-40-6 35661-60-0  
 35737-10-1 35737-15-6 36413-60-2 36653-82-4, 1-Hexadecanol  
 37517-30-9, Acebutolol 41372-20-7 41451-91-6, Erythromycine  
 51264-14-3 51333-22-3 51481-61-9 51781-21-6 53164-05-9,  
 Acemetacin 53716-49-7 54118-66-0 57470-78-7, Celiprolol  
 hydrochloride 57775-29-8, Carazolol 62571-86-2, Captopril  
 63460-06-0 63661-61-0 65571-68-8 68858-20-8 71135-23-4  
 71989-16-7 71989-23-6 71989-28-1 71989-31-6 73724-45-5  
 73731-37-0 75364-47-5 81207-65-0 87980-20-9 91000-69-0  
 91394-66-0 92954-90-0 99103-36-3 104993-76-2 106323-02-8  
 116611-64-4 118477-74-0, Anthron 121343-82-6 125385-14-0,  
 Acyclidine 139369-44-1 139369-45-2 139369-47-4 139369-49-6  
 139369-50-9 139369-51-0 139369-52-1 139390-98-0 139406-80-7  
 139411-94-2 139424-41-2 139465-03-5, Adonit M

(binding of, to anti-dinitrophenol monoclonal antibody, allergic  
 cross-reaction mechanisms in relation to)

L71 ANSWER 18 OF 24 HCAPLUS COPYRIGHT 2002 ACS

1988:575672 Document No. 109:175672 Borohydride, micellar, and exciplex-enhanced dechlorination of chlorobiphenyls. Epling, Gary A.; Florio, Emily M.; Bourque, Andre J.; Qian, Xhi Hong; Stuart, James D. (Dep. Chem., Univ. Connecticut, Storrs, CT, 06268, USA). Environmental Science and Technology, 22(8), 952-6 (English) 1988. CODEN: ESTHAG. ISSN: 0013-936X.


AB The photodechlorination of polychlorinated biphenyls (PCB's) was studied in the presence of NaBH<sub>4</sub>, detergents, and exciplex-forming additives. In a family of 13 representative PCB's, these additives generally led to a dramatically increased rate of photodegrdn. Further, the products of photoreaction in the presence of NaBH<sub>4</sub> are more cleanly the simple dechlorinated aroms., with fewer side **reactions** than obsd. with ordinary photolysis.

IT 9004-95-9, Brij 58

(catalysts, for photochem. dechlorination of chlorinated biphenyls, waste treatment in relation to)

RN 9004-95-9 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)




IT 2437-79-8 35693-99-3, 2,5,2',5'-

Tetrachlorobiphenyl

(photochem. dechlorination of, sodium borohydride- and detergent- and exciplex-enhanced, waste treatment in relation to)


RN 2437-79-8 HCAPLUS

CN 1,1'-Biphenyl, 2,2',4,4'-tetrachloro- (9CI) (CA INDEX NAME)



RN 35693-99-3 HCAPLUS

CN 1,1'-Biphenyl, 2,2',5,5'-tetrachloro- (9CI) (CA INDEX NAME)




IT 33979-03-2, 2,4,6,2',4',6'-Hexachlorobiphenyl  
41411-62-5

(photochem. dechlorination of, sodium borohydride- and  
detergent-enhanced, waste treatment in relation to)


RN 33979-03-2 HCAPLUS

CN 1,1'-Biphenyl, 2,2',4,4',6,6'-hexachloro- (9CI) (CA INDEX NAME)



RN 41411-62-5 HCAPLUS

CN 1,1'-Biphenyl, 2,3,3',4,5,6-hexachloro- (9CI) (CA INDEX NAME)




IT 7012-37-5, 2,4,4'-Trichlorobiphenyl 18259-05-7,  
2,3,4,5,6-Pentachlorobiphenyl 38444-81-4,  
2,5,3'-Trichlorobiphenyl

(photochem. dechlorination of, sodium borohydride- and  
exciplex-enhanced, waste treatment in relation to)


RN 7012-37-5 HCAPLUS

CN 1,1'-Biphenyl, 2,4,4'-trichloro- (9CI) (CA INDEX NAME)




RN 18259-05-7 HCAPLUS

CN 1,1'-Biphenyl, 2,3,4,5,6-pentachloro- (9CI) (CA INDEX NAME)



RN 38444-81-4 HCAPLUS

CN 1,1'-Biphenyl, 2,3',5-trichloro- (9CI) (CA INDEX NAME)




IT 33284-52-5, 3,5,3',5'-Tetrachlorobiphenyl

(photochem. dechlorination of, sodium borohydride-enhanced, waste treatment in relation to)


RN 33284-52-5 HCAPLUS

CN 1,1'-Biphenyl, 3,3',5,5'-tetrachloro- (9CI) (CA INDEX NAME)



- CC 60-4 (Waste Treatment and Disposal)
- IT 121-44-8, Triethylamine, uses and miscellaneous 544-40-1, Dibutyl sulfide 577-11-7, Sodium dioctyl sulfosuccinate 626-17-5, 1,3-Dicyanobenzene 1322-36-7, Dodecanethiol 9004-95-9, Brij 58 16940-66-2, Sodium borohydride (catalysts, for photochem. dechlorination of chlorinated biphenyls, waste treatment in relation to)
- IT 92-52-4D, Biphenyl, chloro derivs. 2050-68-2, 4,4'-Dichlorobiphenyl 2051-60-7, 2-Chlorobiphenyl 2051-62-9, 4-Chlorobiphenyl 2437-79-8 35693-99-3, 2,5,2',5'-Tetrachlorobiphenyl (photochem. dechlorination of, sodium borohydride- and detergent-enhanced, waste treatment in relation to)
- IT 33979-03-2, 2,4,6,2',4',6'-Hexachlorobiphenyl 41411-62-5 (photochem. dechlorination of, sodium borohydride- and detergent-enhanced, waste treatment in relation to)
- IT 2051-61-8, 3-Chlorobiphenyl 7012-37-5, 2,4,4'-Trichlorobiphenyl 18259-05-7, 2,3,4,5,6-Pentachlorobiphenyl 38444-81-4, 2,5,3'-Trichlorobiphenyl (photochem. dechlorination of, sodium borohydride- and exciplex-enhanced, waste treatment in relation to)
- IT 33284-52-5, 3,5,3',5'-Tetrachlorobiphenyl 34883-41-5, 3,5-Dichlorobiphenyl (photochem. dechlorination of, sodium borohydride-enhanced, waste treatment in relation to)
- L71 ANSWER 19 OF 24 HCAPLUS COPYRIGHT 2002 ACS
- 1987:182155 Document No. 106:182155 Process for the decomposition and decontamination of organic substances and halogenated toxic materials. Tundo, Pietro (Sea Marconi Technologies S.p.A., Italy). U.S. US 4632742 A 19861230, 13 pp. Cont.-in-part of U.S. Ser. No. 517,781, abandoned. (English). CODEN: USXXAM. APPLICATION: US 1985-711404 19850313. PRIORITY: IT 1983-19992 19830310; US 1983-517781 19830727.
- AB Haloorg. compds. are **reacted** in the absence of O with a reagent consisting of .gtoreq.1 of a polyethylene glycol, Nixolen, and alc., or a polyhydroxy compd.; a weak base; and an oxidizing agent or radical source. The method is used to decontaminate and recover contaminated industrial oil or decontaminate soil and surfaces contaminated by .gtoreq.1 polyhalogenated org. compd. Transformer oil contg. 20,000 ppm PCB was heated at 85.degree., carbowax 6000 2.0, K2CO3 0.5, and Na2O2 0.2 g were added with stirring for 3 h, the mixt. was cooled, and a clear portion was dild. with n-hexane and analyzed. The residual PCB concn. was 95 ppm.
- IT 50-29-3, biological studies 58-89-9, Hexachlorocyclohexane (decompn. of, in oils and soils, compns. for)
- RN 50-29-3 HCAPLUS
- CN Benzene, 1,1'-(2,2,2-trichloroethylidene)bis[4-chloro- (9CI) (CA


INDEX NAME)



RN 58-89-9 HCAPLUS

CN Cyclohexane, 1,2,3,4,5,6-hexachloro-, (1.alpha.,2.alpha.,3.beta.,4.alpha.,5.alpha.,6.beta.)- (9CI) (CA INDEX NAME)


Relative stereochemistry.



IT 9002-92-0

(haloorg. contaminant decompn. with compns. contg.)

RN 9002-92-0 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-dodecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)

IC ICM B01J019-08

ICS B01J019-10; B01J019-12

NCL 204158210

CC 60-6 (Waste Treatment and Disposal)

Section cross-reference(s): 19, 51

ST chloroorg contaminated oil **reactive** decontamination;  
 chlorodibenzodioxin contaminated oil treatment; PCB contaminated oil  
 treatment; soil haloorg contaminated treatment compn;  
 chlorodibenzofuran decompn polyethylene glycol compn; Nixolen comp  
 decompn haloorg contaminant; alc comp decompn haloorg contaminant

IT 50-29-3, biological studies 58-89-9,



Hexachlorocyclohexane 71-43-2D, Benzene, chloro derivs.  
92-52-4D, Biphenyl, chloro derivs. 108-95-2D, Phenol, chloro  
derivs. 132-64-9D, Dibenzofuran, chloro derivs. 143-50-0  
262-12-4D, chloro derivs. 1746-01-6, 2,3,7,8-Tetrachlorodibenzo-p-  
dioxin

(decompn. of, in oils and soils, compns. for)

IT 78-67-1 112-60-7, Tetraethylene glycol 124-41-4, Sodium  
methoxide 463-79-6D, Carbonic acid, alkali and alk. earth salts  
497-19-8, Disodium carbonate, uses and miscellaneous 584-08-7,  
Dipotassium carbonate 865-47-4 1304-29-6, Barium peroxide  
1313-60-6, Sodium peroxide 9002-92-0 9003-11-6  
25322-68-3, Carbowax 6000 144-55-8, Sodium bicarbonate, uses and  
miscellaneous

(haloorg. contaminant decompn. with compns. contg.)

L71 ANSWER 20 OF 24 HCAPLUS COPYRIGHT 2002 ACS


1987:121884 Document No. 106:121884 Purification of dry cleaning  
solvents with cation exchange-active silicate filter aids.  
Gruenewaelder, Werner; Schaefer, Margarete; Wichelhaus, Winfried;  
Von Rybinski, Wolfgang (Henkel K.-G.a.A., Fed. Rep. Ger.). Ger.  
Offen. DE 3522932 A1 19870108, 7 pp. (German). CODEN: GWXXBX.  
APPLICATION: DE 1985-3522932 19850627.

AB Used drycleaning solvents are purified with the help of a natural  
and/or synthetic silicate filter aid with a cation exchange capacity  
> 30 mequiv/100 g silicate modified with 5-60% .gtoreq.1  
R1R2R3R4N+X- (R1-R4 = H, alkyl, alkenyl; X- = water sol. acid anion  
with dissocn. const. >10<sup>-5</sup>), and with a settling or cartridge filter  
contg. a modified silicate layer alone or in combination with  
kieselguhr and/or activated charcoal. A stirred reactor  
was charged with 500 g Laponite RD and 10 L H2O, heated to  
60.degree. with formation of a thixotropic gel, 425 g Dehyquart LT  
(lauryltrimethylammonium chloride) added, and the mixt. allowed to  
stand for 30 min after stirring. The org.-modified Hecetorite was  
filtered, washed with 2 L H2O, and dried at 75.degree.. The product  
had a C content of 17.3%, contg. .apprx.27% lauryltrimethylammonium  
chloride. A trichlorotrifluoroethane drycleaning solvent, contg. 5  
g/L sebum fatty acids, and 7 g/L of a cleaning enhancer mixt. contg.  
40% Na dodecylbenzene sulfonate, and K petroleum sulfonate 10,  
iso-PrOH 10, spindle oil 20, and H2O 20 g was pumped through a  
filter aid mixt. contg. 60% activated charcoal, and 40% modified  
silicate. The filtered solvent contained sebum fatty acids 1,  
cleaning enhancer 0, and solids content 1.0 g/L, vs. 4, 5, and 7.5,  
resp., for control filter aid mixt. contg. unmodified Hecetorite.


IT 71-55-6P, 1,1,1-Trichloroethane 79-01-6P,  
Trichloroethylene, preparation  
(purifn. of, quaternary ammonium compd.-modified silicate filter  
aids for)

RN 71-55-6 HCAPLUS


CN Ethane, 1,1,1-trichloro- (8CI, 9CI) (CA INDEX NAME)



RN 79-01-6 HCAPLUS  
 CN Ethene, trichloro- (9CI) (CA INDEX NAME)



IT 9004-95-9, Polyethylene glycol monocetyl ether  
 (removal of, from used drycleaning solvents, quaternary ammonium  
 compd.-modified silicate filter aids for)  
 RN 9004-95-9 HCAPLUS  
 CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI)  
 (CA INDEX NAME)




IC ICM D06L001-10  
 ICS B01J020-18; C01B033-20  
 CC 46-5 (Surface Active Agents and Detergents)  
 Section cross-reference(s): 23  
 IT 71-55-6P, 1,1,1-Trichloroethane 79-01-6P,  
 Trichloroethylene, preparation  
 (purifn. of, quaternary ammonium compd.-modified silicate filter  
 aids for)  
 IT 107-21-1, Ethylene glycol, uses and miscellaneous 111-76-2, Butyl  
 glycol 9004-95-9, Polyethylene glycol monocetyl ether  
 9016-45-9, Polyethylene glycol monononylphenol ether 27323-41-7  
 (removal of, from used drycleaning solvents, quaternary ammonium  
 compd.-modified silicate filter aids for)

L71 ANSWER 21 OF 24 HCAPLUS COPYRIGHT 2002 ACS  
 1986:63765 Document No. 104:63765 Peroxide removal from organic  
 solvents and vegetable oils. Shertzer, Howard G.; Tabor, M. Wilson  
 (Med. Cent., Univ. Cincinnati, Cincinnati, OH, 45267-0056, USA).  
 Journal of Environmental Science and Health, Part A: Environmental  
 Science and Engineering, A20(8), 845-55 (English) 1985. CODEN:  
 JESEDU. ISSN: 0360-1226.  
 AB Straight-forward procedures for the testing and removal and(or)

prevention of formation of peroxides in org. solvents and vegetable oils were investigated. Total peroxides were assayed by measuring  $\text{Fe}(\text{SCN})_3$  formed by  $\text{Fe}^{2+}$  oxidn. and **reaction** with KSCN. Most solvents and reagents, as received from the manufacturer, had low levels of peroxides. After opening the container, peroxides were formed rapidly in many solvents. Techniques for the complete retardation of peroxide formation in fresh solvents or reagents are described. For peroxides already present in org. solvents or vegetable oils, a novel technique (employing  $\text{FeSO}_4$  or ferrous-Dowex) for easy removal without introducing contaminants is described.

IT 67-66-3, biological studies 56939-70-9  
 (peroxides of, removal in relation to)  
 RN 67-66-3 HCAPLUS  
 CN Methane, trichloro- (9CI) (CA INDEX NAME)



RN 56939-70-9 HCAPLUS  
 CC 4-3 (Toxicology)  
 IT 50-00-0, biological studies 56-23-5, biological studies 57-55-6,  
 biological studies 60-29-7, biological studies 64-17-5,  
 biological studies 67-56-1, biological studies 67-63-0,  
 biological studies 67-64-1, biological studies 67-66-3,  
 biological studies 67-68-5, biological studies 71-23-8,  
 biological studies 71-36-3, biological studies 71-41-0,  
 biological studies 71-43-2, biological studies 75-05-8,  
 biological studies 75-09-2, biological studies 75-65-0,  
 biological studies 98-06-6 108-86-1, biological studies  
 108-90-7, biological studies 109-66-0, biological studies  
 109-86-4 109-99-9, biological studies 110-54-3, biological  
 studies 110-82-7, biological studies 111-65-9, biological  
 studies 112-40-3 123-51-3 123-54-6, biological studies  
 141-78-6, biological studies 538-23-8 1330-20-7, biological  
 studies 7732-18-5, biological studies 9002-93-1 9004-98-2  
 9005-65-6 9005-66-7 56939-70-9  
 (peroxides of, removal in relation to)

L71 ANSWER 22 OF 24 HCAPLUS COPYRIGHT 2002 ACS

1981:79364 Document No. 94:79364 The principles of enzyme  
 stabilization. VI. Catalysis by water-soluble enzymes entrapped  
 into reversed micelles of surfactants in organic solvents.  
 Martinek, Karel; Levashov, A. V.; Klyachko, N. L.; Pantin, V. I.;  
 Berezin, I. V. (Chem. Dep., Moscow State Univ., Moscow, 117234,  
 USSR). Biochimica et Biophysica Acta, 657(1), 277-94 (English)  
 1981. CODEN: BBACAQ. ISSN: 0006-3002.

AB The possibility of stabilizing water-sol. enzymes against the  
 inactivating action of org. solvents by means of surfactants was  
 studied. Several enzymes [chymotrypsin (EC 3.4.21.1), trypsin (EC


3.4.21.4), inorg. pyrophosphatase (EC 3.6.1.1), peroxidase (EC 1.11.1.7), lactate dehydrogenase (EC 1.1.1.27), and pyruvate kinase (EC 2.7.1.40)] were used to demonstrate that enzymes can be entrapped into reverse micelles formed by surfactants (Aerosol OT, cetyltrimethylammonium bromide, Brij 56) in an org. solvent (benzene, CHCl<sub>3</sub>, octane, cyclohexane). The enzymes solubilized in this way retain their catalytic activity and substrate specificity. A kinetic theory was proposed that describes enzymic **reactions** occurring in a micelle-solvent pseudobiphasic system. In terms of this theory, an explanation is given for the exptl. dependence of the Michaelis-Menten equation parameters on the concns. of the components of a medium (water, org. solvent, surfactant) and also on the combination of the signs of charges in the substrate mol. and at the interphase (++,+,-,--). The results obtained may prove important for applications of enzymes in org. synthesis and for studying the state and role of water in the structure of biomembranes and active sites of enzymes.

IT 9004-95-9

(reverse micelles of, in org. solvents, enzyme entrapment in)

RN 9004-95-9 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)




IT 67-66-3, biological studies

(surfactant reverse micelles in, enzyme entrapment in)

RN 67-66-3 HCAPLUS

CN Methane, trichloro- (9CI) (CA INDEX NAME)



CC 7-13 (Enzymes)

IT 5800-34-0

(**reaction** of, with chymotrypsin entrapped in reverse micelles, kinetics of)

IT 911-76-2

(**reaction** of, with trypsin entrapped in reverse micelles, kinetics of)

IT 57-09-0 577-11-7 9004-95-9

(reverse micelles of, in org. solvents, enzyme entrapment in)

IT 67-66-3, biological studies 71-43-2, biological studies

110-82-7, biological studies 111-65-9, biological studies

(surfactant reverse micelles in, enzyme entrapment in)

L71 ANSWER 23 OF 24 HCAPLUS COPYRIGHT 2002 ACS


1980:491082 Document No. 93:91082 Catalysis by water-soluble enzymes incorporated into reversed surfactant micelles in nonaqueous solvents. Levashov, A. V.; Klyachko, N. L.; Pantin, V. I.; Khmel'nitskii, Yu. L.; Martinek, K. (M. V. Lomonosov State Univ., Moscow, USSR). Bioorganicheskaya Khimiya, 6(6), 929-43 (Russian) 1980. CODEN: BIKHD7. ISSN: 0132-3423.

AB Water-sol. enzyme stabilization with surfactants against the inactivation induced by org. solvents was studied. .alpha.-Chymotrypsin, trypsin, pyrophosphatase, lactate dehydrogenase, pyruvate kinase, and peroxidase were incorporated into reversed micelles formed by surfactants (Aerosol OT, cetyltrimethylammonium bromide, Brij-56) in org. solvents (benzene, CHCl<sub>3</sub>, octane, cyclohexane). Solubilized enzymes retained their catalytic activity and substrate specificity. A kinetic theory of enzyme **reactions** in the pseudo-biphasic micelle-solvent system was proposed. Anal. of the obsd. dependence of K<sub>m</sub> values on concns. of the medium components (water, org. solvent, surfactant) and on the ratio of the charge signs both in the substrate mol. and at the interface (++, +-, --) was carried out. The results obtained can be of value for enzyme application in org. synthesis and for elucidating the state and role of water incorporated into biomembranes and enzyme active centers.

IT 67-66-3, biological studies  
(enzymes incorporated in reversed surfactant micelles in, activity of)

RN 67-66-3 HCAPLUS

CN Methane, trichloro- (9CI) (CA INDEX NAME)




IT 9004-95-9

(reverse micelles of, in org. solvents, enzymes incorporated in, activity of)

RN 9004-95-9 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI)  
(CA INDEX NAME)



CC 7-13 (Enzymes)

IT 67-66-3, biological studies 71-43-2, biological studies  
110-82-7, biological studies 111-65-9, biological studies

(enzymes incorporated in reversed surfactant micelles in, activity of)

IT 57-09-0 577-11-7 9004-95-9

(reverse micelles of, in org. solvents, enzymes incorporated in, activity of)

L71 ANSWER 24 OF 24 HCAPLUS COPYRIGHT 2002 ACS

1967:18062 Document No. 66:18062 The elimination of pesticide residues on apricots by washing before canning. Viel, Guy; Hascoet, M.; Dubroca, G. (Lab. Phytopharm., Inst. Natl. Rech. Agron, Versailles, Fr.). Phytiatr.-Phytopharm., 15(1), 41-8 (French) 1966. CODEN: PHPHA6.


AB Several detergents were tried for the elimination of S on apricots. The effect of the washings was assessed by the use of 35S. About 60 treated apricots were soaked in 10 l. H<sub>2</sub>O or soln., brushed mech. for 2 min., and rinsed for 30 sec. by spraying. Half of the lot was analyzed immediately, the other half was canned to study eventual corrosion control. The washing treatments consisted of (A) H<sub>2</sub>O (control), (B) H<sub>2</sub>O and 0.002% dodecylbenzenesulfonate, (C) H<sub>2</sub>O and 0.0125% oxyethylenated oleocetyl alc. (I) (D) H<sub>2</sub>O and 0.05% oxyethylenated sucrose tallow acid esters (II) and (E) H<sub>2</sub>O with a mixt. of 0.03% I and 0.02% II. Unwashed apricots contained 120-30 mg. S/kg. Compared to the control treatment, the S content was reduced to 5.1% by B, to 17% by C, to 18% by D, and to 14% by E. Treatment D was preferred, as the detergent was biodegradable. Treatment C was also tried on apricots treated with DDT or with carbaryl. A H<sub>2</sub>O washing reduced DDT content by 18% and carbaryl by 75%, but C reduced DDT by 28.5% and carbaryl by 83.5%.

IT 9004-95-9 32054-77-6

(pesticide removal from apricots by polyethylene glycol 9-octadecenyl ether and)

RN 9004-95-9 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-hexadecyl-.omega.-hydroxy- (9CI) (CA INDEX NAME)




RN 32054-77-6 HCAPLUS

IT 50-29-3, biological studies

(residues of, removal from apricots by washing)

RN 50-29-3 HCAPLUS

CN Benzene, 1,1'-(2,2,2-trichloroethylidene)bis[4-chloro- (9CI) (CA INDEX NAME)



- CC 17 (Foods)  
IT Tallow  
    (fatty acids of, esters with sucrose, **reaction** products  
    with ethylene oxide, pesticide removal from apricots by)  
IT Sucrose, esters with tallow fatty acids  
    (**reaction** products with ethylene oxide, pesticide  
    removal from apricots by)  
IT Ethylene oxide  
    Ethylene oxide  
    (**reaction** products with tallow fatty acid esters with  
    sucrose, pesticide removal from apricots by)  
IT 9004-95-9 32054-77-6  
    (pesticide removal from apricots by polyethylene glycol  
    9-octadecenyl ether and)  
IT 50-29-3, biological studies 63-25-2  
    (residues of, removal from apricots by washing)